

Datasheet: MCA883 BATCH NUMBER 1603

Description:	MOUSE ANTI HUMAN CD62E/CD62P
Specificity:	CD62E/CD62P
Other names:	E-SELECTIN/P-SELECTIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	1.2B6
lsotype:	lgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

rad-antibodies.com/proto	cols.			
	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			10ug/ml
Immunohistology - Frozen				1ug/ml - 10ug/ml
(1)	_			rug/mi - roug/mi
Immunohistology - Paraffin		•		
ELISA	-			10ug/ml
Immunoprecipitation	-			
Western Blotting (2)	-			
Where this antibody has necessarily exclude its us a guide only. It is recomm system using appropriate (1) The opitane recognit	se in such nended th e negative	n procedu at the use /positive o	res. Suggested workin er titrates the antibody controls.	g dilutions are given a for use in their own
(1)The epitope recognis	•		• •	
formaldehyde fixation a	and tissu	e proces	sing. Bio-Rad recom	mends the use of
acetone fixation for from	zen secti	ons.		
(2)Clone 1.2B6 recognia	zes huma	n CD62E	CD62P only under r	on-reducing
			CD62P only under r	on-reducing

conditions.

Target Species	Human
Species Cross Reactivity	Reacts with: Pig N.B. Antibody reactivity and working conditions may vary between species. Cross
	reactivity is derived from testing within our laboratories, peer-reviewed publications or

	personal communications from the originators. Please refer to references indicated for further information.			
Product Form	Purified IgG - liquid			
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant			
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	0.09% Sodium Azide			
Carrier Free	Yes			
Approx. Protein Concentrations	IgG concentration 1 mg/ml			
Immunogen	Human E-Selectin (ELAM-1).			
External Database Links	UniProt:P16581Related reagentsP16109Related reagentsEntrez Gene:6401SELERelated reagents6403SELPRelated reagents			
Synonyms	ELAM1, GMRP, GRMP			
RRID	AB_2301666			
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.			
Specificity	Mouse anti Human CD62E/CD62P antibody, clone 1.2B6 recognizes the human CD62E and CD62P cell surface antigens.			
	Although initially thought to recognize only human CD62E, more recent data (<u>Goda et 2003</u>) shows that Mouse anti Human CD62E/CD62P antibody, clone 1.2B6 also recognizes human CD62P, binding to a common epitope shared by these members of selectin family.			
	Clone 1.2B6 reacts with porcine E-selectin (CD62E) but not with porcine P-selectin (Stocker <i>et al.</i> 2000).			
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.			

Histology Positive Control Tissue	Human Tonsil
References	 Wellicome, S.M. <i>et al.</i> (1990) A monoclonal antibody that detects a novel antigen on endothelial cells that is induced by tumor necrosis factor, IL-1, or lipopolysaccharide. J Immunol. 144 (7): 2558-65. Thornhill, M.H. & Haskard, D.O. (1990) IL-4 regulates endothelial cell activation by IL-1, tumor necrosis factor, or IFN-gamma. J Immunol. 145 (3): 865-72. Kyan-Aung, U. <i>et al.</i> (1991) Endothelial leukocyte adhesion molecule-1 and intercellular adhesion molecule-1 mediate the adhesion of eosinophils to endothelial cells <i>in vitro</i> and are expressed by endothelium in allergic cutaneous inflammation <i>in vivo</i>. J Immunol. 146 (2): 521-8. Keelan, E.T. <i>et al.</i> (1994) Characterization of E-selectin expression in vivo with use of a radiolabeled monoclonal antibody. Am J Physiol. 266 (1 Pt 2): H278-90. Goda, K. <i>et al.</i> (1999) Characterization of an apparently conserved epitope in E- and P-selectin identified by dual-specific monoclonal antibodies. Eur J Immunol. 29 (5): 1551-60. Urquhart, P. <i>et al.</i> (2007) Carbon monoxide-releasing molecules modulate leukocyte- endothelial interactions under flow. J Pharmacol Exp Ther 321: 656-662. Gómez del Moral, M. <i>et al.</i> (1999) African swine fever virus infection induces tumor necrosis factor alpha production: implications in pathogenesis. J Virol. 73: 2173-80. Vallée, I. <i>et al.</i> (2001) African swine fever virus infection of porcine aortic endothelial cells leads to inhibition of inflammatory responses, activation of the thrombotic state, and apoptosis. J Virol. 75: 10372-82. Stocker, C.J. <i>et al.</i> (2017) Accelerated resolution of inflammation underlies sex differences in inflammatory responses in humans. J Clin Invest. 127 (1): 169-182.
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA883 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Ar	nti Mouse IgG (STAR12)	E	<u>RPE</u>				
Goat Ant	i Mouse IgG IgA IgM (STAR	37) <u>F</u>	IRP				
Goat Ant	i Mouse IgG (STAR76)	E	RPE				
Rabbit Ar	nti Mouse IgG (STAR13)	Ŀ	<u>IRP</u>				
Goat Ant	Goat Anti Mouse IgG (STAR70) <u>FITC</u>						
Goat Ant	Goat Anti Mouse IgG (H/L) (STAR117) <u>Alk. Phos., DyLight®488</u> , <u>DyLight®550</u> ,						
		<u>[</u>	0yLight®650, DyLight®68	0, DyLight®80	<u>)0</u> ,		
		E	ITC, HRP				
Rabbit Ar	nti Mouse IgG (STAR9)	E	<u>ITC</u>				
Goat Anti Mouse IgG (STAR77)			HRP				
Goat Anti Mouse IgG (Fc) (STAR120) <u>FITC</u> , <u>HRP</u>							
Recomn	nended Negative Control	ls					
MOUSE IgG1 NEGATIVE CONTROL (MCA928)							
North & South	Tel: +1 800 265 7376 Wo	rldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21		
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50		
	Email: antibody_sales_us@bio-rad.com	1	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com		
To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets							

'M369085:200529'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint