

Datasheet: MCA883

**BATCH NUMBER 160959**

<b>Description:</b>	MOUSE ANTI HUMAN CD62E/CD62P
<b>Specificity:</b>	CD62E/CD62P
<b>Other names:</b>	E-SELECTIN/P-SELECTIN
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	1.2B6
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.2 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			10ug/ml
Immunohistology - Frozen (1)	▪			1ug/ml - 10ug/ml
Immunohistology - Paraffin		▪		
ELISA	▪			10ug/ml
Immunoprecipitation	▪			
Western Blotting (2)	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.**

**(2)Clone 1.2B6 recognizes human CD62E/CD62P only under non-reducing conditions.**

<b>Target Species</b>	Human
-----------------------	-------

<b>Species Cross Reactivity</b>	<p>Reacts with: Pig</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or</p>
---------------------------------	---

personal communications from the originators. Please refer to references indicated for further information.

---

<b>Product Form</b>	Purified IgG - liquid
---------------------	-----------------------

---

<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
--------------------	---

---

<b>Buffer Solution</b>	Phosphate buffered saline
------------------------	---------------------------

---

<b>Preservative Stabilisers</b>	0.09% Sodium Azide
---------------------------------	--------------------

---

<b>Carrier Free</b>	Yes
---------------------	-----

---

<b>Approx. Protein Concentrations</b>	IgG concentration 1 mg/ml
---------------------------------------	---------------------------

---

<b>Immunogen</b>	Human E-Selectin (ELAM-1).
------------------	----------------------------

---

<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P16581</a> <a href="#">Related reagents</a> <a href="#">P16109</a> <a href="#">Related reagents</a>  <b>Entrez Gene:</b> <a href="#">6401</a> SELE <a href="#">Related reagents</a> <a href="#">6403</a> SELP <a href="#">Related reagents</a>
--------------------------------	--

---

<b>Synonyms</b>	ELAM1, GMRP, GRMP
-----------------	-------------------

---

<b>RRID</b>	AB_2301666
-------------	------------

---

<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.
------------------------	---

---

<b>Specificity</b>	<p><b>Mouse anti Human CD62E/CD62P antibody, clone 1.2B6</b> recognizes the human CD62E and CD62P cell surface antigens.</p> <p>Although initially thought to recognize only human CD62E, more recent data (<a href="#">Goda et al. 2003</a>) shows that Mouse anti Human CD62E/CD62P antibody, clone 1.2B6 also recognizes human CD62P, binding to a common epitope shared by these members of the selectin family.</p> <p>Clone 1.2B6 reacts with porcine E-selectin (CD62E) but not with porcine P-selectin (<a href="#">Stocker et al. 2000</a>).</p>
--------------------	---

---

<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
-----------------------	---

---

**Histology Positive  
Control Tissue**

Human Tonsil

---

**References**

1. Thornhill, M.H. & Haskard, D.O. (1990) IL-4 regulates endothelial cell activation by IL-1, tumor necrosis factor, or IFN-gamma. [J Immunol. 145 \(3\): 865-72.](#)
2. Keelan, E.T. *et al.* (1994) Characterization of E-selectin expression *in vivo* with use of a radiolabeled monoclonal antibody. [Am J Physiol. 266 \(1 Pt 2\): H278-90.](#)
3. Kyan-Aung, U. *et al.* (1991) Endothelial leukocyte adhesion molecule-1 and intercellular adhesion molecule-1 mediate the adhesion of eosinophils to endothelial cells *in vitro* and are expressed by endothelium in allergic cutaneous inflammation *in vivo*. [J Immunol. 146 \(2\): 521-8.](#)
4. Gómez del Moral, M. *et al.* (1999) African swine fever virus infection induces tumor necrosis factor alpha production: implications in pathogenesis. [J Virol. 73: 2173-80.](#)
5. Goda, K. *et al.* (1999) Characterization of an apparently conserved epitope in E- and P-selectin identified by dual-specific monoclonal antibodies. [Eur J Immunol. 29 \(5\): 1551-60.](#)
6. Stocker, C.J. *et al.* (2000) TNF-alpha, IL-4, and IFN-gamma regulate differential expression of P- and E-selectin expression by porcine aortic endothelial cells. [J Immunol. 164: 3309-15.](#)
7. Vallée, I. *et al.* (2001) African swine fever virus infection of porcine aortic endothelial cells leads to inhibition of inflammatory responses, activation of the thrombotic state, and apoptosis. [J Virol. 75: 10372-82.](#)
8. Urquhart, P. *et al.* (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. [J Pharmacol Exp Ther 321: 656-62.](#)

---

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety  
Information**

Material Safety Datasheet documentation #10040 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA883>  
10040

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

- |   |  |
|---|--|
| Goat Anti Mouse IgG (STAR77...)         | <a href="#">HRP</a>                              |
| Rabbit Anti Mouse IgG (STAR12...)       | <a href="#">RPE</a>                              |
| Goat Anti Mouse IgG IgA IgM (STAR87...) | <a href="#">Alk. Phos.</a> , <a href="#">HRP</a> |
| Goat Anti Mouse IgG (STAR76...)         | <a href="#">RPE</a>                              |

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)  
Goat Anti Mouse IgG (STAR70...) [FITC](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),  
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),  
[FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

## Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M381520:210512'

Printed on 12 Aug 2023

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)