

Datasheet: MCA883

Description:	MOUSE ANTI HUMAN CD62E/CD62P
Specificity:	CD62E/CD62P
Other names:	E-SELECTIN/P-SELECTIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	1.2B6
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			10ug/ml
Immunohistology - Frozen (1)	▪			1ug/ml - 10ug/ml
Immunohistology - Paraffin		▪		
ELISA	▪			10ug/ml
Immunoprecipitation	▪			
Western Blotting (2)	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

(2)Clone 1.2B6 recognizes human CD62E/CD62P only under non-reducing conditions.

Target Species	Human
-----------------------	-------

Species Cross Reactivity	<p>Reacts with: Pig</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for</p>
---------------------------------	--

further information.

Product Form	Purified IgG - liquid
---------------------	-----------------------

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
--------------------	---

Buffer Solution	Phosphate buffered saline
------------------------	---------------------------

Preservative Stabilisers	0.09% Sodium Azide
---------------------------------	--------------------

Carrier Free	Yes
---------------------	-----

Approx. Protein Concentrations	IgG concentration 1 mg/ml
---------------------------------------	---------------------------

Immunogen	Human E-Selectin (ELAM-1).
------------------	----------------------------

External Database Links	<p>UniProt:</p> <p>P16581 Related reagents</p> <p>P16109 Related reagents</p> <p>Entrez Gene:</p> <p>6401 SELE Related reagents</p> <p>6403 SELP Related reagents</p>
--------------------------------	---

Synonyms	ELAM1, GMRP, GRMP
-----------------	-------------------

RRID	AB_2301666
-------------	------------

Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.
------------------------	---

Specificity	<p>Mouse anti Human CD62E/CD62P antibody, clone 1.2B6 recognizes the human CD62E and CD62P cell surface antigens.</p> <p>Although initially thought to recognize only human CD62E, more recent data (Goda et al. 2003) shows that Mouse anti Human CD62E/CD62P antibody, clone 1.2B6 also recognizes human CD62P, binding to a common epitope shared by these members of the selectin family.</p> <p>Clone 1.2B6 reacts with porcine E-selectin (CD62E) but not with porcine P-selectin (Stocker et al. 2000).</p>
--------------------	---

Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
-----------------------	---

Histology Positive	Human Tonsil
---------------------------	--------------

Control Tissue

References

1. Thornhill, M.H. & Haskard, D.O. (1990) IL-4 regulates endothelial cell activation by IL-1, tumor necrosis factor, or IFN-gamma. [J Immunol. 145 \(3\): 865-72.](#)
2. Keelan, E.T. *et al.* (1994) Characterization of E-selectin expression *in vivo* with use of a radiolabeled monoclonal antibody. [Am J Physiol. 266 \(1 Pt 2\): H278-90.](#)
3. Kyan-Aung, U. *et al.* (1991) Endothelial leukocyte adhesion molecule-1 and intercellular adhesion molecule-1 mediate the adhesion of eosinophils to endothelial cells *in vitro* and are expressed by endothelium in allergic cutaneous inflammation *in vivo*. [J Immunol. 146 \(2\): 521-8.](#)
4. Gómez del Moral, M. *et al.* (1999) African swine fever virus infection induces tumor necrosis factor alpha production: implications in pathogenesis. [J Virol. 73: 2173-80.](#)
5. Goda, K. *et al.* (1999) Characterization of an apparently conserved epitope in E- and P-selectin identified by dual-specific monoclonal antibodies. [Eur J Immunol. 29 \(5\): 1551-60.](#)
6. Stocker, C.J. *et al.* (2000) TNF-alpha, IL-4, and IFN-gamma regulate differential expression of P- and E-selectin expression by porcine aortic endothelial cells. [J Immunol. 164: 3309-15.](#)
7. Vallée, I. *et al.* (2001) African swine fever virus infection of porcine aortic endothelial cells leads to inhibition of inflammatory responses, activation of the thrombotic state, and apoptosis. [J Virol. 75: 10372-82.](#)
8. Urquhart, P. *et al.* (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. [J Pharmacol Exp Ther 321: 656-62.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA883>
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

- | | |
|---|--|
| Goat Anti Mouse IgG (STAR77...) | HRP |
| Rabbit Anti Mouse IgG (STAR12...) | RPE |
| Goat Anti Mouse IgG IgA IgM (STAR87...) | Alk. Phos. , HRP |
| Goat Anti Mouse IgG (STAR76...) | RPE |
| Goat Anti Mouse IgG (Fc) (STAR120...) | FITC , HRP |

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M381520:210512'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)