

Datasheet: MCA883FT BATCH NUMBER 166571

Description:	MOUSE ANTI HUMAN CD62E/CD62P:FITC
Specificity:	CD62E/CD62P
Other names:	E-SELECTIN/P-SELECTIN
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	1.2B6
Isotype:	lgG1
Quantity:	25 µg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes No	Not Determined	Suggested Dilution		
	Flow Cytometry	-		Neat		
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.					
Target Species	Human					
Species Cross	Reacts with: Pig					
Reactivity	N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.					
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid.					
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)			
	FITC	490	525			
Preparation	Purified IgG prepared supernatant.	by affinity chromatog	graphy on Protein A fror	n tissue culture		

Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin			
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml			
Immunogen	Human E-Selectin (ELAM-1).			
External Database Links	UniProt: P16581 Related reagents P16109 Related reagents Entrez Gene: 6401 SELE Related reagents 6403 SELP Related reagents			
Synonyms	ELAM1, GMRP, GRMP			
RRID	AB_1102250			
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.			
Specificity	Mouse anti Human CD62E/CD62P antibody, clone 1.2B6 recognizes the human CD62E and CD62P cell surface antigens.			
	Although initially thought to recognize only human CD62E, more recent data (Goda <i>et al.</i> 2003) shows that Mouse anti Human CD62E/CD62P antibody, clone 1.2B6 also recognizes human CD62P, binding to a common epitope shared by these members of the selectin family.			
	Clone 1.2B6 reacts with porcine E-selectin (CD62E) but not with porcine P-selectin (<u>Stocker <i>et al.</i> 2000</u>).			
Flow Cytometry	Use 10ul of the suggested working dilution to label 1 x 10^6 cells in 100ul.			
References	 Thornhill, M.H. & Haskard, D.O. (1990) IL-4 regulates endothelial cell activation by IL-1, tumor necrosis factor, or IFN-gamma. <u>J Immunol. 145 (3): 865-72.</u> Keelan, E.T. <i>et al.</i> (1994) Characterization of E-selectin expression <i>in vivo</i> with use of a radiolabeled monoclonal antibody. <u>Am J Physiol. 266 (1 Pt 2): H278-90.</u> Kyan-Aung, U. <i>et al.</i> (1991) Endothelial leukocyte adhesion molecule-1 and intercellular adhesion molecule-1 mediate the adhesion of eosinophils to endothelial cells <i>in vitro</i> and are expressed by endothelium in allergic cutaneous inflammation <i>in vivo</i>. <u>J Immunol. 146 (2): 521-8.</u> Gómez del Moral, M. <i>et al.</i> (1999) African swine fever virus infection induces tumor 			

	necrosis factor alpha production: implications in pathogenesis. <u>J Virol. 73: 2173-80.</u> 5. Goda, K. <i>et al.</i> (1999) Characterization of an apparently conserved epitope in E- and
	P-selectin identified by dual-specific monoclonal antibodies. <u>Eur J Immunol. 29 (5):</u> <u>1551-60.</u>
	 6. Stocker, C.J. <i>et al.</i> (2000) TNF-alpha, IL-4, and IFN-gamma regulate differential expression of P- and E-selectin expression by porcine aortic endothelial cells. <u>J Immunol.</u> 164: 3309-15. 7. Vallée, I. <i>et al.</i> (2001) African swine fever virus infection of porcine aortic endothelial cells leads to inhibition of inflammatory responses, activation of the thrombotic state, and apoptosis. <u>J Virol. 75: 10372-82.</u> 8. Urquhart, P. <i>et al.</i> (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. <u>J Pharmacol Exp Ther 321: 656-62.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA883FT 10041
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@	bio-rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385346:210513'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint