

Datasheet: MCA883T

Description:	MOUSE ANTI HUMAN CD62E/CD62P
Specificity:	CD62E/CD62P
Other names:	E-SELECTIN/P-SELECTIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	1.2B6
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			10ug/ml
Immunohistology - Frozen (1)	▪			1ug/ml - 10ug/ml
Immunohistology - Paraffin		▪		
ELISA	▪			10ug/ml
Immunoprecipitation	▪			
Western Blotting (2)	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)**The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.**

(2)**Clone 1.2B6 recognizes human CD62E/CD62P only under non-reducing conditions.**

Target Species	Human
Species Cross Reactivity	<p>Reacts with: Pig</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for</p>

further information.

Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Human E-Selectin (ELAM-1).
External Database Links	UniProt: P16581 Related reagents P16109 Related reagents Entrez Gene: 6401 SELE Related reagents 6403 SELP Related reagents
Synonyms	ELAM1, GMRP, GRMP
RRID	AB_1102249
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.
Specificity	<p>Mouse anti Human CD62E/CD62P antibody, clone 1.2B6 recognizes the human CD62E and CD62P cell surface antigens.</p> <p>Although initially thought to recognize only human CD62E, more recent data (Goda <i>et al.</i> 2003) shows that Mouse anti Human CD62E/CD62P antibody, clone 1.2B6 also recognizes human CD62P, binding to a common epitope shared by these members of the selectin family.</p> <p>Clone 1.2B6 reacts with porcine E-selectin (CD62E) but not with porcine P-selectin (Stocker <i>et al.</i> 2000).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10^6 cells in 100ul.
Histology Positive	Human Tonsil

References

1. Thornhill, M.H. & Haskard, D.O. (1990) IL-4 regulates endothelial cell activation by IL-1, tumor necrosis factor, or IFN-gamma. [J Immunol. 145 \(3\): 865-72.](#)
2. Keelan, E.T. *et al.* (1994) Characterization of E-selectin expression *in vivo* with use of a radiolabeled monoclonal antibody. [Am J Physiol. 266 \(1 Pt 2\): H278-90.](#)
3. Kyan-Aung, U. *et al.* (1991) Endothelial leukocyte adhesion molecule-1 and intercellular adhesion molecule-1 mediate the adhesion of eosinophils to endothelial cells *in vitro* and are expressed by endothelium in allergic cutaneous inflammation *in vivo*. [J Immunol. 146 \(2\): 521-8.](#)
4. Gómez del Moral, M. *et al.* (1999) African swine fever virus infection induces tumor necrosis factor alpha production: implications in pathogenesis. [J Virol. 73: 2173-80.](#)
5. Goda, K. *et al.* (1999) Characterization of an apparently conserved epitope in E- and P-selectin identified by dual-specific monoclonal antibodies. [Eur J Immunol. 29 \(5\): 1551-60.](#)
6. Stocker, C.J. *et al.* (2000) TNF-alpha, IL-4, and IFN-gamma regulate differential expression of P- and E-selectin expression by porcine aortic endothelial cells. [J Immunol. 164: 3309-15.](#)
7. Vallée, I. *et al.* (2001) African swine fever virus infection of porcine aortic endothelial cells leads to inhibition of inflammatory responses, activation of the thrombotic state, and apoptosis. [J Virol. 75: 10372-82.](#)
8. Rathod, K.S. *et al.* (2017) Accelerated resolution of inflammation underlies sex differences in inflammatory responses in humans. [J Clin Invest. 127 \(1\): 169-82.](#)
9. Urquhart, P. *et al.* (2007) Carbon monoxide-releasing molecules modulate leukocyte-endothelial interactions under flow. [J Pharmacol Exp Ther 321: 656-62.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA883T>
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)

Goat Anti Mouse IgG (STAR76...) [RPE](#)

Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG (STAR77...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M381901:210512'

Printed on 19 Aug 2024