

Datasheet: MCA884A647

**BATCH NUMBER 1701**

<b>Description:</b>	RAT ANTI MOUSE CD169:Alexa Fluor® 647
<b>Specificity:</b>	CD169
<b>Other names:</b>	SIALOADHESIN
<b>Format:</b>	ALEXA FLUOR® 647
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	3D6.112
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	100 TESTS/1ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG conjugated to Alexa Fluor 647 - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.05 mg/ml
<b>Immunogen</b>	Purified murine sialoadhesin.

**External Database****Links****UniProt:**[Q62230](#)[Related reagents](#)**Entrez Gene:**[20612](#)

Siglec1

[Related reagents](#)**Synonyms**

Sa, Sn

**Fusion Partners**

Spleen cells from an immunised AO rat were fused with the cells of the Y3 rat myeloma cell line.

**Specificity**

**Rat anti Mouse CD169 antibody, clone 3D6.112** recognizes mouse CD169 also known as sialoadhesin, Sheep erythrocyte receptor or Siglec-1. CD169 is a 1695 amino acid, ~180 kDa single pass, type 1 transmembrane glycoprotein containing a single [Ig-like V-type](#) domain and sixteen [Ig-like C2-type](#) domains. CD169 is a macrophage restricted receptor, preferentially binding to alpha 2,3 linked sialic acid residues ([Crocker et al. 1991](#)) and is expressed on stromal macrophages in many tissues, particularly in lymph nodes, bone marrow and on marginal metallophilic macrophages in the spleen ([Morris et al. 1991](#)).

CD169 has been implicated in a number of roles including cell-cell interactions with lymphocytes ([van den Berg et al. 1992](#)) and granulocytes ([Crocker et al. 1995](#)). CD169 expressing macrophages have also been suggested to play a role in host resistance to lymphoma metastasis ([Umansky et al. 1996](#)). In pigs CD169 has also been identified as a macrophage restricted receptor for porcine reproductive and respiratory syndrome virus ([Delputte et al. 2007](#)). CD169 expressing macrophages have also been implicated in the regulation of autoimmune disease progression through their interaction with regulatory T cells via CD169 ([Wu et al. 2009](#)). CD169 has also been shown to play a critical role in the recognition and elimination of invasive sialylated microorganisms including *Campylobacter jejuni* ([Klass et al. 2012](#)) and group B Streptococcus ([Chang et al. 2014](#)).

The functional activity of rat anti mouse CD169 antibody, clone 3D6.112, its ability to inhibit binding of red blood cells to CD169 can be considerably enhanced by derivitization of the antibody with polyethylene glycol ([Ducreux et al. 2008](#)).

**Flow Cytometry**

Use 10ul of the suggested working dilution to label  $1 \times 10^6$  cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

**References**

1. Crocker, P.R. *et al.* (1991) Purification and properties of sialoadhesin, a sialic acid-binding receptor of murine tissue macrophages. [EMBO J. 10 \(7\): 1661-9.](#)
2. Sancho-Pelluz, J. *et al.* (2008) Sialoadhesin expression in intact degenerating retinas and following transplantation. [Invest Ophthalmol Vis Sci. 49: 5602-10.](#)
3. Barral, P. *et al.* (2010) CD169(+) macrophages present lipid antigens to mediate early activation of iNKT cells in lymph nodes. [Nat Immunol. 11: 303-12.](#)
4. Chtanova, T. *et al.* (2008) Dynamics of neutrophil migration in lymph nodes during infection. [Immunity. 29: 487-96.](#)

5. Hsu, K.M. *et al.* (2009) Murine cytomegalovirus displays selective infection of cells within hours after systemic administration. [J Gen Virol. 90:33-43.](#)
6. Iannacone M (2010) Subcapsular sinus macrophages prevent CNS invasion on peripheral infection with a neurotropic virus. [Nature. 465: 1079-83.](#)
7. Idoyaga, J. *et al.* (2009) Antibody to Langerin/CD207 localizes large numbers of CD8alpha+ dendritic cells to the marginal zone of mouse spleen. [Proc Natl Acad Sci U S A. 106: 1524-9.](#)
8. Taylor, P.R. *et al.* (2008) Development of a specific system for targeting protein to metallophilic macrophages. [Proc Natl Acad Sci U S A. 101: 1963-8.](#)
9. Chang, Y.C. *et al.* (2014) Role of macrophage sialoadhesin in host defense against the sialylated pathogen group B Streptococcus. [J Mol Med \(Berl\). 92\(9\):951-9.](#)
10. Lin, H.H. *et al.* (2005) The macrophage F4/80 receptor is required for the induction of antigen-specific efferent regulatory T cells in peripheral tolerance. [J Exp Med. 201: 1615-25.](#)
11. Phillips, R. *et al.* (2010) Innate killing of *Leishmania donovani* by macrophages of the splenic marginal zone requires IRF-7. [PLoS Pathog. 6\(3\):e1000813.](#)
12. Hashimoto, D. *et al.* (2011) Pretransplant CSF-1 therapy expands recipient macrophages and ameliorates GVHD after allogeneic hematopoietic cell transplantation. [J Exp Med. 208: 1069-82.](#)
13. Anderson, K.L. *et al.* (1999) PU.1 and the granulocyte- and macrophage colony-stimulating factor receptors play distinct roles in late-stage myeloid cell differentiation. [Blood. 94: 2310-8.](#)
14. Albacker, L.A. *et al.* (2010) TIM-4, a receptor for phosphatidylserine, controls adaptive immunity by regulating the removal of antigen-specific T cells. [J Immunol. 185: 6839-49.](#)
15. Anthony, R.M. *et al.* (2008) Identification of a receptor required for the anti-inflammatory activity of IVIG. [Proc Natl Acad Sci U S A. 105: 19571-8.](#)
16. Chow, A. *et al.* (2011) Bone marrow CD169+ macrophages promote the retention of hematopoietic stem and progenitor cells in the mesenchymal stem cell niche. [J Exp Med. 208: 261-71.](#)
17. Hemmi, H. *et al.* (2009) A new triggering receptor expressed on myeloid cells (Trem) family member, Trem-like 4, binds to dead cells and is a DNAX activation protein 12-linked marker for subsets of mouse macrophages and dendritic cells. [J Immunol. 182: 1278-86.](#)
18. Hemmi, H. *et al.* (2012) Trem14, an Ig superfamily member, mediates presentation of several antigens to T cells in vivo, including protective immunity to HER2 protein. [J Immunol. 188: 1147-55.](#)
19. Huang, Q.Q. *et al.* (2010) FLIP: a novel regulator of macrophage differentiation and granulocyte homeostasis. [Blood. 116: 4968-77.](#)
20. Lu, M. and Munford, R.S. (2011) The transport and inactivation kinetics of bacterial lipopolysaccharide influence its immunological potency *in vivo*. [J Immunol. 187: 3314-20.](#)
21. Krücken, J. *et al.* (2005) Massive destruction of malaria-parasitized red blood cells despite spleen closure. [Infect Immun. 73: 6390-8.](#)
22. Vagaja, N.N. *et al.* (2012) Changes in murine hyalocytes are valuable early indicators of ocular disease. [Invest Ophthalmol Vis Sci. 53: 1445-51.](#)
23. Barnes, Y.C. *et al.* (1999) Sialylation of the sialic acid binding lectin sialoadhesin regulates its ability to mediate cell adhesion. [Blood. 93: 1245-52.](#)
24. Chen, W.C. *et al.* (2012) Antigen delivery to macrophages using liposomal nanoparticles targeting sialoadhesin/CD169. [PLoS One. 7e39039.](#)

25. Mansour, A. *et al.* (2012) Osteoclasts promote the formation of hematopoietic stem cell niches in the bone marrow. [J Exp Med. 209: 537-49.](#)
26. Park, M.H. *et al.* (2015) Neuropeptide Y regulates the hematopoietic stem cell microenvironment and prevents nerve injury in the bone marrow. [EMBO J. 34 \(12\): 1648-60.](#)
27. Asai, H. *et al.* (2015) Depletion of microglia and inhibition of exosome synthesis halt tau propagation. [Nat Neurosci. 18 \(11\): 1584-93.](#)
28. Farrell, H.E. *et al.* (2015) Lymph Node Macrophages Restrict Murine Cytomegalovirus Dissemination. [J Virol. 89 \(14\): 7147-58.](#)
29. Xu, H.C. *et al.* (2015) Deficiency of the B cell-activating factor receptor results in limited CD169+ macrophage function during viral infection. [J Virol. 89 \(9\): 4748-59.](#)
30. Gao, L. *et al.* (2015) Infiltration of circulating myeloid cells through CD95L contributes to neurodegeneration in mice. [J Exp Med. 212 \(4\): 469-80.](#)
31. McCabe, A. *et al.* (2015) Macrophage-Lineage Cells Negatively Regulate the Hematopoietic Stem Cell Pool in Response to Interferon Gamma at Steady State and During Infection. [Stem Cells. 33 \(7\): 2294-305.](#)
32. Cantisani, R. *et al.* (2015) Vaccine adjuvant MF59 promotes retention of unprocessed antigen in lymph node macrophage compartments and follicular dendritic cells. [J Immunol. 194 \(4\): 1717-25.](#)
33. Prokopec, K.E. *et al.* (2016) Marginal Zone Macrophages Regulate Antigen Transport by B Cells to the Follicle in the Spleen via CD21. [J Immunol. 197 \(6\): 2063-8.](#)
34. Li, Z. *et al.* (2016) The Macrophage-depleting Agent Clodronate Promotes Durable Hematopoietic Chimerism and Donor-specific Skin Allograft Tolerance in Mice. [Sci Rep. 6: 22143.](#)
35. Farrell, H.E. *et al.* (2016) Type 1 Interferons and NK Cells Limit Murine Cytomegalovirus Escape from the Lymph Node Subcapsular Sinus. [PLoS Pathog. 12 \(12\): e1006069.](#)
36. Perrotta, M. *et al.* (2018) Deoxycorticosterone acetate-salt hypertension activates placental growth factor in the spleen to couple sympathetic drive and immune system activation. [Cardiovasc Res. 114 \(3\): 456-67.](#)
37. Urata, S. *et al.* (2018) BST-2 controls T cell proliferation and exhaustion by shaping the early distribution of a persistent viral infection. [PLoS Pathog. 14 \(7\): e1007172.](#)
38. Peng, Y. (2018) B cell responses to apoptotic cells in MFG-E8-/- mice. [PLoS One. 13 \(10\): e0205172.](#)
39. Tay, M.H.D. *et al.* (2019) Halted Lymphocyte Egress via Efferent Lymph Contributes to Lymph Node Hypertrophy During Hypercholesterolemia. [Front Immunol. 10: 575.](#)
40. Ding, X. *et al.* (2019) Panicle-Shaped Sympathetic Architecture in the Spleen Parenchyma Modulates Antibacterial Innate Immunity. [Cell Rep. 27 \(13\): 3799-3807.e3.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.  
 Storage in frost-free freezers is not recommended.  
 This product should be stored undiluted. This product is photosensitive and should be protected from light.  
 Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Acknowledgements** This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchased product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or [outlicensing@thermofisher.com](mailto:outlicensing@thermofisher.com)

---

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA884A647>  
10041

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA1212A647\)](#)

### Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M369091:200529'

**Printed on 19 Jun 2025**

---

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)