

Datasheet: MCA88AMO

| | |
|----------------------|---|
| Description: | MOUSE ANTI HUMAN CD45RA:Amethyst Orange |
| Specificity: | CD45RA |
| Format: | Amethyst Orange |
| Product Type: | Monoclonal Antibody |
| Clone: | F8-11-13 |
| Isotype: | IgG1 |
| Quantity: | 0.1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | Neat - 1/10 |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Rhesus Monkey

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to Amethyst Orange - liquid

Max Ex/Em

| Fluorophore | Excitation Max (nm) | Emission Max (nm) |
|-----------------|---------------------|-------------------|
| Amethyst Orange | 405 | 540 |

Preparation

Purified IgG prepared by affinity chromatography on protein A from tissue culture supernatant

Buffer Solution

Phosphate buffered saline

| | |
|---------------------------------------|---|
| Preservative Stabilisers | 0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin |
| Approx. Protein Concentrations | IgG concentration 0.1 mg/ml |
| Immunogen | Human T lymphocytes |
| External Database Links | <p>UniProt: P08575 Related reagents</p> <p>Entrez Gene: 5788 PTPRC Related reagents</p> |
| Synonyms | CD45 |
| Fusion Partners | Spleen cells from immunized BALB/c mouse were fused with cells of the NS-1 mouse myeloma cell line |
| Specificity | <p>Mouse anti Human CD45RA antibody, clone F8-11-13 recognizes the human CD45RA cell surface antigen.</p> <p>CD45 is a complex molecule existing in a number of isoforms. Antibodies recognizing a common epitope on all of these isoforms are termed CD45 whilst those recognizing only individual isoforms are termed CD45RA or CD45RO etc.</p> <p>Mouse anti Human CD45RA antibody, clone F8-11-13 reacts with the high molecular weight form of the leucocyte common antigen (LCA) expressed by B lymphocytes and a subset of T lymphocytes.</p> <p>Mouse anti Human CD45RA antibody, clone F8-11-13 is routinely tested in flow cytometry on human peripheral blood lymphocytes.</p> |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul |
| References | <ol style="list-style-type: none"> 1. Dalchau, R. & Fabre, J.W. (1981) Identification with a monoclonal antibody of a predominantly B lymphocyte-specific determinant of the human leukocyte common antigen. Evidence for structural and possible functional diversity of the human leukocyte common molecule. J Exp Med. 153 (4): 753-65. 2. O'Grady, J.T. <i>et al.</i> (1994) CD40 expression in Hodgkin's disease. Am J Pathol. 144 (1): 21-6. 3. Franitza, S. <i>et al.</i> (2002) TGF-beta1 enhances SDF-1alpha-induced chemotaxis and homing of naive T cells by up-regulating CXCR4 expression and downstream cytoskeletal effector molecules. Eur J Immunol. 32 (1): 193-202. 4. Leigh, J.E. <i>et al.</i> (2006) Characterization of the immune status of CD8+ T cells in oral lesions of human immunodeficiency virus-infected persons with oropharyngeal Candidiasis. Clin Vaccine Immunol. 13 (6): 678-83. |

5. Hunt, L. *et al.* (2016) T cell subsets: an immunological biomarker to predict progression to clinical arthritis in ACPA-positive individuals. [Ann Rheum Dis. 75 \(10\): 1884-9.](#)
6. Ibane, J.A. *et al.* (2012) The major CD8 T cell effector memory subset in the normal and *Chlamydia trachomatis*-infected human endocervix is low in perforin. [BMC Immunol. 13: 66.](#)
7. Ponchel, F. *et al.* (2014) An immunological biomarker to predict MTX response in early RA. [Ann Rheum Dis. 73 \(11\): 2047-53.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA88AMO>
10041

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Amethyst Orange \(MCA928AMO\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M385059:210513'

Printed on 18 Jan 2024