

Datasheet: MCA890T BATCH NUMBER 154511

| Description: | MOUSE ANTI HUMAN CYTOKERATIN 14 | | |
|---------------|---------------------------------|--|--|
| Specificity: | CYTOKERATIN 14 | | |
| Format: | Purified | | |
| Product Type: | Monoclonal Antibody | | |
| Clone: | LL002 | | |
| Isotype: | IgG3 | | |
| Quantity: | 20 μg | | |
| | | | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry (1) | • | | | 1/100 |
| Immunohistology - Frozen | • | | | |
| Immunohistology - Paraffin (2) | - | | | 1/200 |
| ELISA | | | • | |
| Immunoprecipitation | | | • | |
| Immunofluorescence | | | | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

- (1)Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm[™] (Product Code <u>BUF09</u>) for this purpose.
- (2)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

| Target Species | Human |
|----------------|--|
| Species Cross | Reacts with: Elephant, Dog, Pig, Lion |
| Reactivity | N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. |

| Product Form | Purified IgG - liquid | | |
|--------------------------------------|--|--|--|
| Preparation | Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant | | |
| Buffer Solution | Phosphate buffered saline | | |
| Preservative Stabilisers | 0.09% Sodium Azide | | |
| Carrier Free | Yes | | |
| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml | | |
| Immunogen | Last 15 C-terminal residues of human cytokeratin 14 conjugated to thyroglobulin. | | |
| External Database Links | UniProt: P02533 Related reagents Entrez Gene: 3861 KRT14 Related reagents | | |
| RRID | AB_323916 | | |
| Specificity | Mouse anti Human Cytokeratin 14 antibody, clone LL002 recognizes cytokeratin 14, a type I intermediate filament, expressed by stratifying epithelial cells and can be used to distinguish these cell types from simple epithelial cells, which do not express cytokeratin 14. | | |
| | Mouse anti cytokeratin 14, clone LL002 has been reported to be suitable for use in Western blotting (<u>Alam et al. 2011</u>) | | |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul. | | |
| Histology Positive Control Tissue | Skin | | |
| References | Purkis, P.E. <i>et al.</i> (1990) Antibody markers of basal cells in complex epithelia. <u>J Cell Sci. 97 (Pt 1): 39-50.</u> Lane, E.B. & Alexander, C.M. (1990) Use of keratin antibodies in tumor diagnosis. <u>Semin Cancer Biol. 1 (3): 165-79.</u> Wetzels, R.H. <i>et al.</i> (1989) Detection of basement membrane components and basal cell keratin 14 in noninvasive and invasive carcinomas of the breast. <u>Am J Pathol. 134 (3): 571-9.</u> Moll, R. <i>et al.</i> (1982) The catalog of human cytokeratins: patterns of expression in normal epithelia, tumors and cultured cells. <u>Cell. 31 (1): 11-24.</u> Richardson, G.D. <i>et al.</i> (2004) CD133, a novel marker for human prostatic epithelial | | |

- stem cells. J Cell Sci. 117 (Pt 16): 3539-45.
- 6. Holliday, D. *et al.* (2009) Novel multicellular organotypic models of normal and malignant breast: tools for dissecting the role of the microenvironment in breast cancer progression. <u>Breast Cancer Res. 11: R3</u>
- 7. Eastman, R. .Jr. *et al.* (2010) Fibroblast growth factor-10 signals development of von Brunn's nests in the exstrophic bladder. <u>Am J Physiol Renal Physiol.299:F1094-110.</u>
- 8. Stumpf, P and Welsch, U. (2004) Secretory and defensive functions of the duct system of the lactating mammary gland of the African elephant (*Loxodonta africana*, Proboscidea) Zoomorphology 123:155-67
- 9. Alam H *et al.* (2011) Loss of keratins 8 and 18 leads to alterations in $\alpha6\beta4$ -integrinmediated signalling and decreased neoplastic progression in an oral-tumour-derived cell line. J Cell Sci. 124 (Pt 12): 2096-106.
- 10. Clark, S.E. *et al.* (2011) Molecular subtyping of DCIS: heterogeneity of breast cancer reflected in pre-invasive disease. Br J Cancer. 104: 120-7.
- 11. Caceres, S. *et al.* (2015) Establishment and Characterization of a New Cell Line of Canine Inflammatory Mammary Cancer: IPC-366. PLoS One. 10 (3): e0122277.
- 12. Takahashi, C. *et al.* (2010) Newly established cell lines from mouse oral epithelium regenerate teeth when combined with dental mesenchyme. <u>In Vitro Cell Dev Biol Anim.</u> 46: 457-68.
- 13. Faustino, A.M. & Dias Pereira, P. (2007) A salivary malignant myoepithelioma in a dog. Vet J. 173 (1): 223-6.
- 14. Collins, A.T. *et al*et al. (2005) Prospective identification of tumorigenic prostate cancer stem cells. Cancer Res. 65: 10946-51.
- 15. Varley, C.L. *et al.* (2004) Activation of peroxisome proliferator-activated receptor-gamma reverses squamous metaplasia and induces transitional differentiation in normal human urothelial cells. Am J Pathol. 164: 1789-98.
- 16. Abdeen, S.K. *et al.* (2011) Wwox inactivation enhances mammary tumorigenesis. Oncogene. 30: 3900-6.
- 17. Matos, A.J. *et al.* (2006) Detection of lymph node micrometastases in malignant mammary tumours in dogs by cytokeratin immunostaining. Vet Rec. 158: 626-30.
- 18. Turner, A. *et al.* (2011) Transplantation of autologous differentiated urothelium in an experimental model of composite cystoplasty. Eur Urol. 59: 447-54.
- 19. Munz, B. *et al.* (1999) Overexpression of activin A in the skin of transgenic mice reveals new activities of activin in epidermal morphogenesis, dermal fibrosis and wound repair. <u>EMBO J. 18: 5205-15.</u>
- 20. Mwase, M. *et al.* (2013) Cutaneous Squamous Cell Carcinoma presenting as a Wound with Discharging Sinus Tracts in a Wild African Lion (*Panthera leo*). <u>J Comp Pathol. pii:</u> S0021-9975(13)00106-0.
- 21. Honda, K. & Tomooka, Y. (2016) Nerve-independent and ectopically additional induction of taste buds in organ culture of fetal tongues. *In Vitro* Cell Dev Biol Anim. 52 (9): 911-9.
- 22. Hustler, A. *et al.* (2018) Differential transcription factor expression by human epithelial cells of buccal and urothelial derivation. <u>Exp Cell Res. 369 (2): 284-294.</u>
- 23. Ogihara, K. & Madarame, H. (2020) Pancreatic adenosquamous carcinoma with invasion to the spleen in a cat. J Vet Med Sci. 82 (9): 1395-9.

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

| Guarantee | 12 months from date of despatch |
|----------------------------------|--|
| Health And Safety Information | Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA890T 10040 |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) **RPE**

Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

Goat Anti Mouse IgG (STAR76...) **RPE** Goat Anti Mouse IgG (STAR70...) **FITC**

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Goat Anti Mouse IgG (STAR77...) **HRP**

Rabbit Anti Mouse IgG (STAR9...) **FITC**

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) **HRP**

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M372015:200617'

Printed on 30 Jul 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint