

Datasheet: MCA907

BATCH NUMBER 158254

Description:	MOUSE ANTI HUMAN CD106
Specificity:	CD106
Other names:	VCAM-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	1.G11B1
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation	▪			
Western Blotting (1)	▪			
Immunofluorescence	▪			
Functional Assays (2)	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) This product has been reported to work in non-reducing conditions

(2) This product contains sodium azide, removal by dialysis is recommended prior to use in functional assays.

Target Species	Human
-----------------------	-------

Species Cross Reactivity	<p>Reacts with: Pig, Rhesus Monkey</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for</p>
---------------------------------	---

further information.

Product Form	Purified IgG - liquid
---------------------	-----------------------

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
--------------------	---

Buffer Solution	Phosphate buffered saline
------------------------	---------------------------

Preservative Stabilisers	0.09% Sodium Azide
---------------------------------	--------------------

Carrier Free	Yes
---------------------	-----

Approx. Protein Concentrations	IgG concentration 1 mg/ml
---------------------------------------	---------------------------

External Database Links	UniProt: P19320 Related reagents Entrez Gene: 7412 VCAM1 Related reagents
--------------------------------	--

Synonyms	L1CAM
-----------------	-------

RRID	AB_321992
-------------	-----------

Specificity	<p>Mouse anti Human CD106 antibody, clone 1.G11B1 recognizes human VCAM-1, a ~110 kDa molecule whose ligand is VLA4. The antigen is expressed on activated endothelial cells and on some tissue macrophages, bone marrow fibroblasts and myoblasts.</p> <p>Mouse anti Human CD106 antibody, clone 1.G11B1 inhibits cellular adhesion mediated by VCAM-1 (Patel 1998).</p>
--------------------	--

References	<ol style="list-style-type: none">1. Thornhill, M.H. <i>et al.</i> (1991) Tumor necrosis factor combines with IL-4 or IFN-gamma to selectively enhance endothelial cell adhesiveness for T cells. The contribution of vascular cell adhesion molecule-1-dependent and -independent binding mechanisms. J Immunol. 146 (2): 592-8.2. Rosenman, S.J. <i>et al.</i> (1995) Cytokine-induced expression of vascular cell adhesion molecule-1 (VCAM-1) by astrocytes and astrocytoma cell lines. J Immunol. 154 (4): 1888-99.3. Peterson, M.D. <i>et al.</i> (2005) Monocyte-induced endothelial calcium signaling mediates early xenogeneic endothelial activation. Am J Transplant 5 (2): 237-47.4. Reparon-Schuijt, C.C. <i>et al.</i> (2000) Regulation of synovial B cell survival in rheumatoid arthritis by vascular cell adhesion molecule 1 (CD106) expressed on fibroblast-like synoviocytes. Arthritis Rheum. 43 (5): 1115-21.5. Faure, J.P. <i>et al.</i> (2002) Polyethylene glycol reduces early and long-term cold ischemia-
-------------------	--

- reperfusion and renal medulla injury. [J Pharmacol Exp Ther. 302 \(3\): 861-70.](#)
6. Ruschulte, H. *et al.* (2011) Adrenoceptor stimulation does not affect ICAM-1 and VCAM-1 expression *in vitro*. [BMC Res Notes. 4: 40.](#)
 7. Old, E.A. *et al.* (2014) Monocytes expressing CX3CR1 orchestrate the development of vincristine-induced pain. [J Clin Invest. 124 \(5\): 2023-36.](#)
 8. Cabeza, N. *et al.* (2004) Surface expression of collagen receptor Fc receptor-gamma/glycoprotein VI is enhanced on platelets in type 2 diabetes and mediates release of CD40 ligand and activation of endothelial cells. [Diabetes 53: 2117-21.](#)
 9. Corvaisier, M. *et al.* (2005) V gamma 9V delta 2 T cell response to colon carcinoma cells. [J Immunol. 175: 5481-8.](#)
 10. Dunoyer-Geindre, S. *et al.* (2004) Aspirin inhibits endothelial cell activation induced by antiphospholipid antibodies. [J Thromb Haemost. 2: 1176-81.](#)
 11. Hauet, T. *et al.* (2002) Polyethylene glycol reduces the inflammatory injury due to cold ischemia/reperfusion in autotransplanted pig kidneys. [Kidney Int. 62: 654-67.](#)
 12. Kindle, L. *et al.* (2005) Human microvascular endothelial cell activation by IL-1 and TNF-alpha stimulates the adhesion and transendothelial migration of circulating human CD14+ monocytes that develop with RANKL into functional osteoclasts. [J Bone Miner Res. 21: 193-206.](#)
 13. Kahler, C.M. (2007) Peripheral infusion of rat bone marrow derived endothelial progenitor cells leads to homing in acute lung injury. [Respir Res. 8: 50.](#)
 14. Norling, L.V. *et al.* (2008) Inhibitory control of endothelial galectin-1 on *in vitro* and *in vivo* lymphocyte trafficking. [FASEB J. 22: 682-90.](#)
 15. Colleselli, D. *et al.* (2006) Inhibition of cyclooxygenase (COX)-2 affects endothelial progenitor cell proliferation. [Exp Cell Res. 312: 2933-41.](#)
 16. Hayashida, K. *et al.* (2000) Rheumatoid arthritis synovial stromal cells inhibit apoptosis and up-regulate Bcl-xL expression by B cells in a CD49/CD29-CD106-dependent mechanism. [J Immunol. 164: 1110-6.](#)
 17. Ali, S. *et al.* (2000) Intercellular cell adhesion molecule-1, vascular cell adhesion molecule-1, and regulated on activation normal T cell expressed and secreted are expressed by human breast carcinoma cells and support eosinophil adhesion and activation. [Am J Pathol. 157: 313-21.](#)
 18. Pelletier M, Girard D. (2005) Interleukin-15 increases neutrophil adhesion onto human respiratory epithelial A549 cells and attracts neutrophils *in vivo*. [Clin Exp Immunol. 141: 315-25.](#)
 19. Murphy, A.J. *et al.* (2013) Anti-inflammatory functions of apolipoprotein a-I and high-density lipoprotein are preserved in trimeric apolipoprotein a-I. [J Pharmacol Exp Ther. 344: 41-9.](#)
 20. Chadderdon, S.M. *et al.* (2014) Proinflammatory endothelial activation detected by molecular imaging in obese nonhuman primates coincides with onset of insulin resistance and progressively increases with duration of insulin resistance. [Circulation. 129 \(4\): 471-8.](#)
 21. Zhang, J. *et al.* (2016) Bone mesenchymal stem cells differentiate into myofibroblasts in the tumor microenvironment. [Oncol Lett. 12 \(1\): 644-50.](#)
 22. Lim, J.L. *et al.* (2016) Protective effects of monomethyl fumarate at the inflamed blood-brain barrier. [Microvasc Res. 105: 61-9.](#)
 23. Holzwarth, C. *et al.* (2010) Low physiologic oxygen tensions reduce proliferation and differentiation of human multipotent mesenchymal stromal cells. [BMC Cell Biol. 11: 11.](#)
 24. Schmidt, M. *et al.* (2016) Methods to Investigate the Role of Toll-Like Receptors in

Allergic Contact Dermatitis. [Methods Mol Biol. 1390: 319-40.](#)

25. May, R.D. *et al.* (2012) Preclinical development of CAT-354, an IL-13 neutralizing antibody, for the treatment of severe uncontrolled asthma. [Br J Pharmacol. 166 \(1\): 177-93.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA907>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M376210:210123'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)