

## Datasheet: MCA907PE

<b>Description:</b>	MOUSE ANTI HUMAN CD106:RPE
<b>Specificity:</b>	CD106
<b>Other names:</b>	VCAM-1
<b>Format:</b>	RPE
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	1.G11B1
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

#### Target Species

Human

#### Species Cross Reactivity

Reacts with: Pig, Rhesus Monkey

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

#### Reconstitution

Reconstitute with 1.0 ml distilled water

Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution

#### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
RPE 488nm laser	496	578

<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Stabilisers</b>	1% Bovine Serum Albumin 5% Sucrose
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P19320</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">7412</a>    VCAM1    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	L1CAM
<b>Specificity</b>	<p><b>Mouse anti Human CD106 antibody, clone 1.G11B1</b> recognizes human VCAM-1, a ~110 kDa molecule whose ligand is VLA4. The antigen is expressed on activated endothelial cells and on some tissue macrophages, bone marrow fibroblasts and myoblasts.</p> <p>Mouse anti Human CD106 antibody, clone 1.G11B1 inhibits cellular adhesion mediated by VCAM-1 (<a href="#">Patel 1998</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul
<b>References</b>	<ol style="list-style-type: none"> <li>1. Thornhill, M.H. <i>et al.</i> (1991) Tumor necrosis factor combines with IL-4 or IFN-gamma to selectively enhance endothelial cell adhesiveness for T cells. The contribution of vascular cell adhesion molecule-1-dependent and -independent binding mechanisms. <a href="#">J Immunol. 146 (2): 592-8.</a></li> <li>2. Rosenman, S.J. <i>et al.</i> (1995) Cytokine-induced expression of vascular cell adhesion molecule-1 (VCAM-1) by astrocytes and astrocytoma cell lines. <a href="#">J Immunol. 154 (4): 1888-99.</a></li> <li>3. Peterson, M.D. <i>et al.</i> (2005) Monocyte-induced endothelial calcium signaling mediates early xenogeneic endothelial activation. <a href="#">Am J Transplant 5 (2): 237-47.</a></li> <li>4. Reparon-Schuijt, C.C. <i>et al.</i> (2000) Regulation of synovial B cell survival in rheumatoid arthritis by vascular cell adhesion molecule 1 (CD106) expressed on fibroblast-like synoviocytes. <a href="#">Arthritis Rheum. 43 (5): 1115-21.</a></li> <li>5. Faure, J.P. <i>et al.</i> (2002) Polyethylene glycol reduces early and long-term cold ischemia-reperfusion and renal medulla injury. <a href="#">J Pharmacol Exp Ther. 302 (3): 861-70.</a></li> <li>6. Ruschulte, H. <i>et al.</i> (2011) Adrenoceptor stimulation does not affect ICAM-1 and VCAM-1 expression <i>in vitro</i>. <a href="#">BMC Res Notes. 4: 40.</a></li> <li>7. Old, E.A. <i>et al.</i> (2014) Monocytes expressing CX3CR1 orchestrate the development of vincristine-induced pain. <a href="#">J Clin Invest. 124 (5): 2023-36.</a></li> <li>8. Cabeza, N. <i>et al.</i> (2004) Surface expression of collagen receptor Fc receptor-gamma/glycoprotein VI is enhanced on platelets in type 2 diabetes and mediates release</li> </ol>

- of CD40 ligand and activation of endothelial cells. [Diabetes 53: 2117-21.](#)
9. Corvaisier, M. *et al.* (2005) V gamma 9V delta 2 T cell response to colon carcinoma cells. [J Immunol. 175: 5481-8.](#)
10. Dunoyer-Geindre, S. *et al.* (2004) Aspirin inhibits endothelial cell activation induced by antiphospholipid antibodies. [J Thromb Haemost. 2: 1176-81.](#)
11. Hauet, T. *et al.* (2002) Polyethylene glycol reduces the inflammatory injury due to cold ischemia/reperfusion in autotransplanted pig kidneys. [Kidney Int. 62: 654-67.](#)
12. Kindle, L. *et al.* (2005) Human microvascular endothelial cell activation by IL-1 and TNF-alpha stimulates the adhesion and transendothelial migration of circulating human CD14+ monocytes that develop with RANKL into functional osteoclasts. [J Bone Miner Res. 21: 193-206.](#)
13. Kahler, C.M. (2007) Peripheral infusion of rat bone marrow derived endothelial progenitor cells leads to homing in acute lung injury. [Respir Res. 8: 50.](#)
14. Norling, L.V. *et al.* (2008) Inhibitory control of endothelial galectin-1 on *in vitro* and *in vivo* lymphocyte trafficking. [FASEB J. 22: 682-90.](#)
15. Colleselli, D. *et al.* (2006) Inhibition of cyclooxygenase (COX)-2 affects endothelial progenitor cell proliferation. [Exp Cell Res. 312: 2933-41.](#)
16. Hayashida, K. *et al.* (2000) Rheumatoid arthritis synovial stromal cells inhibit apoptosis and up-regulate Bcl-xL expression by B cells in a CD49/CD29-CD106-dependent mechanism. [J Immunol. 164: 1110-6.](#)
17. Ali, S. *et al.* (2000) Intercellular cell adhesion molecule-1, vascular cell adhesion molecule-1, and regulated on activation normal T cell expressed and secreted are expressed by human breast carcinoma cells and support eosinophil adhesion and activation. [Am J Pathol. 157: 313-21.](#)
18. Pelletier M, Girard D. (2005) Interleukin-15 increases neutrophil adhesion onto human respiratory epithelial A549 cells and attracts neutrophils *in vivo*. [Clin Exp Immunol. 141: 315-25.](#)
19. Murphy, A.J. *et al.* (2013) Anti-inflammatory functions of apolipoprotein a-I and high-density lipoprotein are preserved in trimeric apolipoprotein a-I. [J Pharmacol Exp Ther. 344: 41-9.](#)
20. Chadderdon, S.M. *et al.* (2014) Proinflammatory endothelial activation detected by molecular imaging in obese nonhuman primates coincides with onset of insulin resistance and progressively increases with duration of insulin resistance. [Circulation. 129 \(4\): 471-8.](#)
21. Zhang, J. *et al.* (2016) Bone mesenchymal stem cells differentiate into myofibroblasts in the tumor microenvironment. [Oncol Lett. 12 \(1\): 644-50.](#)
22. Lim, J.L. *et al.* (2016) Protective effects of monomethyl fumarate at the inflamed blood-brain barrier. [Microvasc Res. 105: 61-9.](#)
23. Holzwarth, C. *et al.* (2010) Low physiologic oxygen tensions reduce proliferation and differentiation of human multipotent mesenchymal stromal cells. [BMC Cell Biol. 11: 11.](#)
24. Schmidt, M. *et al.* (2016) Methods to Investigate the Role of Toll-Like Receptors in Allergic Contact Dermatitis. [Methods Mol Biol. 1390: 319-40.](#)
25. May, R.D. *et al.* (2012) Preclinical development of CAT-354, an IL-13 neutralizing antibody, for the treatment of severe uncontrolled asthma. [Br J Pharmacol. 166 \(1\): 177-93.](#)

---

**Storage**

Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be

protected from light.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #20487 available at: 20487: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M375691:210104'

Printed on 29 Aug 2021