

# Datasheet: MCA90PE BATCH NUMBER 166574

Description:	MOUSE ANTI HUMAN CD90:RPE		
Specificity:	CD90		
Other names:	THY1		
Format:	RPE		
Product Type:	Monoclonal Antibody		
Clone:	F15-42-1		
Isotype:	lgG1		
Quantity:	100 TESTS		

# **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
	Yes No Not Determined Suggested Dilution							
	Flow Cytometry	•		Neat				
	necessarily exclude it	s use in such proce mmended that the u	user titrates the product	ng dilutions are given as				
Target Species	Human							
Species Cross Reactivity	Reacts with: Cynomolgus monkey <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.							
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized							
Reconstitution	Reconstitute in 1 ml distilled water. Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently vortexed after reconstitution and microcentrifuged before use.							
Max Ex/Em	Fluorophore	Excitation Max (nm	) Emission Max (nm)					
	RPE 488nm laser	496	578					

Preparation	Purified IgG prepared by ion exchange chromatography from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	<ul><li>0.09% Sodium Azide</li><li>1% Bovine Serum Albumin</li><li>5% Sucrose</li></ul>
Immunogen	Purified human brain Thy-1.
External Database Links	UniProt:         P04216       Related reagents         Entrez Gene:         7070       THY1         Related reagents
RRID	AB_321888
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.
Specificity	<ul> <li>Mouse anti Human CD90 antibody, clone F15-42-1 recognizes the human CD90 cell surface antigen, a ~25 kDa glycoprotein homologous to rat Thy1. The antigen is expressed by a subset of CD34+ve cells in the bone marrow and by prothymocytes within the thymus. CD90 is also expressed extensively within the brain.</li> <li>Mouse anti Human CD90 antibody, clone F15-42-1 is routinely tested in flow cytometry on the MOLT4 cell line.</li> </ul>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
References	<ol> <li>Daar, A.S. &amp; Fabre, J.W. (1981) Demonstration with monoclonal antibodies of an unusual mononuclear cell infiltrate and loss of normal epithelial membrane antigens in human breast carcinomas. Lancet. 2 (8244): 434-8.</li> <li>Fiegel, H.C. <i>et al.</i> (2004) Stem-like cells in human hepatoblastoma. J Histochem Cytochem. 52 (11): 1495-501.</li> <li>Hagood, J.S. <i>et al.</i> (2005) Loss of fibroblast Thy-1 expression correlates with lung fibrogenesis. Am J Pathol. 167 (2): 365-79.</li> <li>Tome, M. <i>et al.</i> (2007) Calponin is expressed by subpopulations of connective tissue cells but not olfactory ensheathing cells in the neonatal olfactory mucosa. BMC Neurosci. 8: 74.</li> <li>Diaz-Romero, J. <i>et al.</i> (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect bona fide dedifferentiation rather than amplification of progenitor cells. J Cell Physiol. 214: 75-83.</li> <li>Pessina, A. <i>et al.</i> (2010) CD45+/CD133+ positive cells expanded from umbilical cord blood expressing PDX-1 and markers of pluripotency. Cell Biol Int. 34: 783-90.</li> <li>Manochantr, S. <i>et al.</i> (2010) Isolation, characterization and neural differentiation</li> </ol>

potential of amnion derived mesenchymal stem cells. <u>J Med Assoc Thai. 93 Suppl 7:</u> <u>S183-91.</u>

8. Karlsen, T.A. *et al.* (2010) Human primary articular chondrocytes, chondroblasts-like cells, and dedifferentiated chondrocytes: differences in gene, microRNA, and protein expression and phenotype. Tissue Eng Part C Methods. 17: 219-27.

9. Hauser, P.V. *et al.* (2010) Stem cells derived from human amniotic fluid contribute to acute kidney injury recovery. <u>Am J Pathol. 177: 2011-21.</u>

10. Yin, S. *et al.* (2010) Chondrogenic transdifferentiation of human dermal fibroblasts stimulated with cartilage-derived morphogenetic protein 1. <u>Tissue Eng Part A. 16:</u> 1633-43.

11. Gieseke, F. *et al.* (2010) Human multipotent mesenchymal stromal cells use galectin-1 to inhibit immune effector cells. <u>Blood. 116: 3770-9.</u>

12. Holzwarth, C. *et al.* (2010) Low physiologic oxygen tensions reduce proliferation and differentiation of human multipotent mesenchymal stromal cells. <u>BMC Cell Biol. 11:11</u>

13. Meng, J. *et al* (2011) Contribution of human muscle-derived cells to skeletal muscle regeneration in dystrophic host mice. <u>PLoS One. 6: e17454.</u>

14. Cizeau, J. *et al.* (2011) Fusogenics: a recombinant immunotoxin-based screening platform to select internalizing tumor-specific antibody fragments. J Biomol Screen. 16: 90-100.

15. Cox, G. *et al.* (2011) The use of the reamer-irrigator-aspirator to harvest mesenchymal stem cells. J Bone Joint Surg Br. 93: 517-24.

16. Shafaei, H. *et al.* (2011) Effects of human placental serum on proliferation and morphology of human adipose tissue-derived stem cells. <u>Bone Marrow Transplant. 46:</u> 1464-71.

17. Paul, G. *et al.* (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. <u>PLoS One. 7: e35577.</u>

18. Supokawej, A. *et al.* (2013) Cardiogenic and myogenic gene expression in mesenchymal stem cells after 5-azacytidine treatment. Turk J Haematol. 30 (2): 115-21.

19. Escobar, C.H. & Chaparro, O. (2016) Xeno-Free Extraction, Culture, and

Cryopreservation of Human Adipose-Derived Mesenchymal Stem Cells. <u>Stem Cells Transl</u> <u>Med. 5 (3): 358-65.</u>

20. Shinoda, K. *et al.* (2016) Thy1+IL-7+ lymphatic endothelial cells in iBALT provide a survival niche for memory T-helper cells in allergic airway inflammation. <u>Proc Natl Acad</u> <u>Sci U S A. 113 (20): E2842-51.</u>

21. Kamprom, W. *et al.* (2016) Endothelial Progenitor Cell Migration-Enhancing Factors in the Secretome of Placental-Derived Mesenchymal Stem Cells. <u>Stem Cells Int. 2016</u>: <u>2514326</u>.

22. Vaquero, J. *et al.* (2016) An approach to personalized cell therapy in chronic complete paraplegia: The Puerta de Hierro phase I/II clinical trial. <u>Cytotherapy. 18 (8): 1025-36.</u>
23. Zhang, X. *et al.* (2017) Regeneration of hyaline-like cartilage in situ with SOX9 stimulation of bone marrow-derived mesenchymal stem cells. <u>PLoS One. 12 (6):</u> e0180138.

24. Garikipati, V. N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. <u>PLoS One. 13 (2): e0192244.</u>

25. Chaturvedi, C.P. *et al.* (2018) Altered Expression of Hematopoiesis Regulatory Molecules in Lipopolysaccharide-Induced Bone Marrow Mesenchymal Stem Cells of Patients with Aplastic Anemia. <u>Stem Cells Int. 2018: 6901761.</u>

	<ol> <li>Noda, S. <i>et al.</i> (2019) Effect of cell culture density on dental pulp-derived mesenchymal stem cells with reference to osteogenic differentiation. <u>Sci Rep. 9 (1): 5430</u>.</li> <li>Song, H. <i>et al.</i> (2019) MIF/CD74 axis participates in inflammatory activation of Schwann cells following sciatic nerve injury. <u>J Mol Histol. 50 (4): 355-67</u>.</li> <li>Paiboon, N. <i>et al.</i> (2019) Gestational Tissue-Derived Human Mesenchymal Stem Cells Use Distinct Combinations of Bioactive Molecules to Suppress the Proliferation of Human Hepatoblastoma and Colorectal Cancer Cells. <u>Stem Cells Int. 2019</u>: 9748795.</li> <li>Sanjurjo-Rodriguez, C. <i>et al.</i> (2020) Gene Expression Signatures of Synovial Fluid Multipotent Stromal Cells in Advanced Knee Osteoarthritis and Following Knee Joint Distraction. <u>Front Bioeng Biotechnol. 8</u>: 579751.</li> <li>Fujii-Tezuka, R. <i>et al.</i> (2021) Umbilical artery tissue contains p75 neurotrophin receptor-positive pericyte-like cells that possess neurosphere formation capacity and neurogenic differentiation potential. <u>Regen Ther. 16</u>: 1-11.</li> <li>Orikasa, S. <i>et al.</i> (2022) Hypoxia-inducible factor 1α induces osteo/odontoblast differentiation of human dental pulp stem cells via Wnt/β-catenin transcriptional cofactor BCL9. <u>Sci Rep. 12 (1): 682</u>.</li> <li>Sirithammajak, S. <i>et al.</i> (2022) Human Mesenchymal Stem Cells Derived from the Placenta and Chorion Suppress the Proliferation while Enhancing the Migration of Human Breast Cancer Cells. <u>Stem Cells Int. 2022</u>: 4020845.</li> <li>Arenal, Á. <i>et al.</i> (2023) Effects of Cardiac Stem Cell on Postinfarction Arrhythmogenic Substrate. <u>Int J Mol Sci. 23 (24): 16211.</u></li> <li>Kruchen, A. <i>et al.</i> (2023) Epigenetic Modification of Mesenchymal Stromal Cells Derived from Bone Marrow and Embryonal Tumors to Facilitate Immunotherapeutic Approaches in Pediatric Malignancies <u>Current Issues in Molecular Biology. 45 (3)</u>: 2121-35.</li> </ol>
Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C. This product should be stored undiluted.
	' DO NOT FREEZE. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA90PE 20487
Regulatory	For research purposes only

## **Related Products**

## **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

#### **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419607:230616'

#### Printed on 29 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint