

# Datasheet: MCA90SBB700

| Description:  | MOUSE ANTI HUMAN CD90:StarBright Blue 700 |
|---------------|---|
| Specificity:  | CD90                                      |
| Other names:  | THY1                                      |
| Format:       | StarBright Blue 700                       |
| Product Type: | Monoclonal Antibody                       |
| Clone:        | F15-42-1                                  |
| lsotype:      | lgG1                                      |
| Quantity:     | 100 TESTS/0.5ml                           |
|               |   |

### **Product Details**

| Applications                | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . |               |           |                  |                    |  |
|-----------------------------|--|---------------|-----------|------------------|--------------------|--|
|                             |  | Yes           | No        | Not Determined   | Suggested Dilution |  |
|                             | Flow Cytometry   | •             |           |                  | Neat               |  |
|                             | Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.   |               |           |                  |                    |  |
| Target Species              | Human  |               |           |                  |                    |  |
| Species Cross<br>Reactivity | Reacts with: Cynomolgus monkey<br><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross<br>reactivity is derived from testing within our laboratories, peer-reviewed publications or<br>personal communications from the originators. Please refer to references indicated for<br>further information.  |               |           |                  |                    |  |
| Product Form                | Purified IgG conjugated to StarBright Blue 700 - liquid  |               |           |                  |                    |  |
| Max Ex/Em                   | Fluorophore  | Excitation Ma | x (nm) Ei | mission Max (nm) |                    |  |
|                             | StarBright Blue 700  | 473           |           | 703              |                    |  |
| Preparation                 | Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant  |               |           |                  |                    |  |
| Buffer Solution             | Phosphate buffered sa  | aline         |           |                  |                    |  |

| Preservative<br>Stabilisers | 0.09% Sodium Azide (NaN <sub>3</sub> )<br>1% Bovine Serum Albumin<br>0.1% Pluronic F68<br>0.1% PEG 3350<br>0.05% Tween 20  |
|-----------------------------|--|
| Immunogen                   | Purified human brain Thy-1.  |
| External Database<br>Links  | UniProt:<br><u>P04216</u> <u>Related reagents</u><br>Entrez Gene:<br><u>7070</u> THY1 <u>Related reagents</u>  |
| Fusion Partners             | Spleen cells from immunized BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.   |
| Specificity                 | <ul> <li>Mouse anti Human CD90 antibody, clone F15-42-1 recognizes the human CD90 cell surface antigen, a ~25 kDa glycoprotein homologous to rat Thy1. The antigen is expressed by a subset of CD34+ve cells in the bone marrow and by prothymocytes within the thymus. CD90 is also expressed extensively within the brain.</li> <li>Mouse anti Human CD90 antibody, clone F15-42-1 is routinely tested in flow cytometry on the MOLT4 cell line.</li> </ul>  |
| Flow Cytometry              | Use 5µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.  |
| References                  | <ol> <li>Daar, A.S. &amp; Fabre, J.W. (1981) Demonstration with monoclonal antibodies of an<br/>unusual mononuclear cell infiltrate and loss of normal epithelial membrane antigens in<br/>human breast carcinomas. <u>Lancet. 2 (8244): 434-8.</u></li> <li>Fiegel, H.C. <i>et al.</i> (2004) Stem-like cells in human hepatoblastoma. <u>J Histochem</u><br/><u>Cytochem. 52 (11): 1495-501.</u></li> <li>Hagood, J.S. <i>et al.</i> (2005) Loss of fibroblast Thy-1 expression correlates with lung<br/>fibrogenesis. <u>Am J Pathol. 167 (2): 365-79.</u></li> <li>Tome, M. <i>et al.</i> (2007) Calponin is expressed by subpopulations of connective tissue<br/>cells but not olfactory ensheathing cells in the neonatal olfactory mucosa. <u>BMC Neurosci.<br/>8: 74.</u></li> <li>Diaz-Romero, J. <i>et al.</i> (2008) Immunophenotypic changes of human articular<br/>chondrocytes during monolayer culture reflect bona fide dedifferentiation rather than<br/>amplification of progenitor cells. <u>J Cell Physiol. 214: 75-83.</u></li> <li>Pessina, A. <i>et al.</i> (2010) CD45+/CD133+ positive cells expanded from umbilical cord<br/>blood expressing PDX-1 and markers of pluripotency. <u>Cell Biol Int. 34: 783-90.</u></li> <li>Manochantr, S. <i>et al.</i> (2010) Isolation, characterization and neural differentiation<br/>potential of amnion derived mesenchymal stem cells. <u>J Med Assoc Thai. 93 Suppl 7:<br/>S183-91.</u></li> <li>Karlsen, T.A. <i>et al.</i> (2010) Human primary articular chondrocytes, chondroblasts-like</li> </ol> |

cells, and dedifferentiated chondrocytes: differences in gene, microRNA, and protein expression and phenotype. <u>Tissue Eng Part C Methods</u>. <u>17: 219-27</u>.

9. Hauser, P.V. *et al.* (2010) Stem cells derived from human amniotic fluid contribute to acute kidney injury recovery. <u>Am J Pathol. 177: 2011-21.</u>

10. Yin, S. *et al.* (2010) Chondrogenic transdifferentiation of human dermal fibroblasts stimulated with cartilage-derived morphogenetic protein 1. <u>Tissue Eng Part A. 16:</u> <u>1633-43.</u>

11. Gieseke, F. *et al.* (2010) Human multipotent mesenchymal stromal cells use galectin-1 to inhibit immune effector cells. <u>Blood. 116: 3770-9.</u>

Holzwarth, C. *et al.* (2010) Low physiologic oxygen tensions reduce proliferation and differentiation of human multipotent mesenchymal stromal cells. <u>BMC Cell Biol. 11:11</u>
 Meng, J. *et al* (2011) Contribution of human muscle-derived cells to skeletal muscle

regeneration in dystrophic host mice. <u>PLoS One. 6: e17454.</u>
14. Cizeau, J. *et al.* (2011) Fusogenics: a recombinant immunotoxin-based screening platform to select internalizing tumor-specific antibody fragments. <u>J Biomol Screen. 16:</u> 90-100.

15. Cox, G. *et al.* (2011) The use of the reamer-irrigator-aspirator to harvest mesenchymal stem cells. J Bone Joint Surg Br. 93: 517-24.

16. Shafaei, H. *et al.* (2011) Effects of human placental serum on proliferation and morphology of human adipose tissue-derived stem cells. <u>Bone Marrow Transplant. 46:</u> 1464-71.

17. Paul, G. *et al.* (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. <u>PLoS One. 7: e35577.</u>

 Supokawej, A. *et al.* (2013) Cardiogenic and myogenic gene expression in mesenchymal stem cells after 5-azacytidine treatment. <u>Turk J Haematol. 30 (2): 115-21.</u>
 Escobar, C.H. & Chaparro, O. (2016) Xeno-Free Extraction, Culture, and

Cryopreservation of Human Adipose-Derived Mesenchymal Stem Cells. <u>Stem Cells Transl</u> <u>Med. 5 (3): 358-65.</u>

20. Shinoda, K. *et al.* (2016) Thy1+IL-7+ lymphatic endothelial cells in iBALT provide a survival niche for memory T-helper cells in allergic airway inflammation. <u>Proc Natl Acad</u> Sci U S A. 113 (20): E2842-51.

21. Kamprom, W. *et al.* (2016) Endothelial Progenitor Cell Migration-Enhancing Factors in the Secretome of Placental-Derived Mesenchymal Stem Cells. <u>Stem Cells Int. 2016</u>: <u>2514326</u>.

22. Vaquero, J. *et al.* (2016) An approach to personalized cell therapy in chronic complete paraplegia: The Puerta de Hierro phase I/II clinical trial. <u>Cytotherapy. 18 (8): 1025-36.</u>
23. Zhang, X. *et al.* (2017) Regeneration of hyaline-like cartilage in situ with SOX9 stimulation of bone marrow-derived mesenchymal stem cells. <u>PLoS One. 12 (6):</u> e0180138.

24. Garikipati, V. N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. <u>PLoS One. 13 (2): e0192244.</u>

25. Chaturvedi, C.P. *et al.* (2018) Altered Expression of Hematopoiesis Regulatory Molecules in Lipopolysaccharide-Induced Bone Marrow Mesenchymal Stem Cells of Patients with Aplastic Anemia. <u>Stem Cells Int. 2018: 6901761.</u>

26. Noda, S. *et al.* (2019) Effect of cell culture density on dental pulp-derived mesenchymal stem cells with reference to osteogenic differentiation. <u>Sci Rep. 9 (1): 5430.</u>
27. Song, H. *et al.* (2019) MIF/CD74 axis participates in inflammatory activation of

| Schwann cells following sciatic nerve injury. <u>J Mol Histol. 50 (4): 355-67.</u> |  |  |  |  |  |  |
|--|--|--|--|--|--|--|
|  | 28. Paiboon, N. et al. (2019) Gestational Tissue-Derived Human Mesenchymal Stem Cells  |  |  |  |  |  |
|  | Use Distinct Combinations of Bioactive Molecules to Suppress the Proliferation of Huma   |  |  |  |  |  |
|  | Hepatoblastoma and Colorectal Cancer Cells. <u>Stem Cells Int. 2019: 9748795.</u>  |  |  |  |  |  |
|  | 29. Sanjurjo-Rodriguez, C. et al. (2020) Gene Expression Signatures of Synovial Fluid  |  |  |  |  |  |
|  | Multipotent Stromal Cells in Advanced Knee Osteoarthritis and Following Knee Joint   |  |  |  |  |  |
|  | Distraction. Front Bioeng Biotechnol. 8: 579751.   |  |  |  |  |  |
|  | 30. Fujii-Tezuka, R. <i>et al.</i> (2021) Umbilical artery tissue contains p75 neurotrophin  |  |  |  |  |  |
|  | receptor-positive pericyte-like cells that possess neurosphere formation capacity and  |  |  |  |  |  |
|  | neurogenic differentiation potential. Regen Ther. 16: 1-11.  |  |  |  |  |  |
|  | 31. Orikasa, S. <i>et al.</i> (2022) Hypoxia-inducible factor 1α induces osteo/odontoblast   |  |  |  |  |  |
|  | differentiation of human dental pulp stem cells via Wnt/β-catenin transcriptional cofactor   |  |  |  |  |  |
|  | BCL9. Sci Rep. 12 (1): 682.  |  |  |  |  |  |
|  | 32. Sirithammajak, S. <i>et al.</i> (2022) Human Mesenchymal Stem Cells Derived from the   |  |  |  |  |  |
|  | Placenta and Chorion Suppress the Proliferation while Enhancing the Migration of Human   |  |  |  |  |  |
|  | Breast Cancer Cells. <u>Stem Cells Int. 2022: 4020845.</u>   |  |  |  |  |  |
|  | 33. Arenal, Á. <i>et al.</i> (2022) Effects of Cardiac Stem Cell on Postinfarction Arrhythmogenic  |  |  |  |  |  |
|  |  |  |  |  |  |  |
|  | Substrate. Int J Mol Sci. 23 (24): 16211.<br>34 Kruchen A. et al. (2023) Epigenetic Modification of Mesenchymal Stromal Cells  |  |  |  |  |  |
|  | 34. Kruchen, A. <i>et al.</i> (2023) Epigenetic Modification of Mesenchymal Stromal Cells  |  |  |  |  |  |
|  | Derived from Bone Marrow and Embryonal Tumors to Facilitate Immunotherapeutic<br>Approaches in Pediatric Malignancies. <u>Curr Issues Mol Biol. 45 (3): 2121-35.</u> |  |  |  |  |  |
|  |  |  |  |  |  |  |
|  | 35. Payet, M. <i>et al.</i> (2023) Inflammatory Mesenchymal Stem Cells Express Abundant  |  |  |  |  |  |
|  | Membrane-Bound and Soluble Forms of C-Type Lectin-like CD248. Int J Mol Sci. 24 (11  |  |  |  |  |  |
|  | <u>9546.</u>   |  |  |  |  |  |
|  | 36. Tiraihi, T. <i>et al.</i> (2023) A Sequential Culturing System for Generating Epithelial-Like  |  |  |  |  |  |
|  | Stem Cells from Human Mesenchymal Stem Cells Derived from Adipose Tissue Cell  |  |  |  |  |  |
|  | <u>Tissue Biol. 17 (6): 639-52.</u>  |  |  |  |  |  |
|  | 37. Tripathy, N.K. et al. (2018) Cardiomyogenic Heterogeneity of Clonal Subpopulations of  |  |  |  |  |  |
|  | Human Bone Marrow Mesenchymal Stem Cells. <u>J Stem Cells Regen Med. 14 (1): 27-33.</u>  |  |  |  |  |  |
|  | 38. Altaie, A. et al. (2022) Device-Based Enrichment of Knee Joint Synovial Cells to Drive   |  |  |  |  |  |
|  | MSC Chondrogenesis Without Prior Culture Expansion In Vitro: A Step Closer to 1-Stage  |  |  |  |  |  |
|  | Orthopaedic Procedures. Am J Sports Med. 50 (1): 152-161.  |  |  |  |  |  |
| 0.4  |  |  |  |  |  |  |
| Storage  | Store at +4°C. DO NOT FREEZE.  |  |  |  |  |  |
|  | This product should be stored undiluted.   |  |  |  |  |  |
| Guarantee  | 12 months from date of despatch  |  |  |  |  |  |
|  |  |  |  |  |  |  |
| Acknowledgements   | This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign   |  |  |  |  |  |
|  | counterparts   |  |  |  |  |  |
| Health And Safety  | Material Safety Datasheet documentation #20471 available at:   |  |  |  |  |  |
| Information  | https://www.bio-rad-antibodies.com/SDS/MCA90SBB700   |  |  |  |  |  |
|  | 20471  |  |  |  |  |  |
|  |  |  |  |  |  |  |
| Regulatory   | For research purposes only   |  |  |  |  |  |
|  |  |  |  |  |  |  |

# **Related Products**

#### HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

| North & South | Tel: +1 800 265 7376                 | Worldwide | Tel: +44 (0)1865 852 700             | Europe | Tel: +49 (0) 89 8090 95 21           |
|---------------|--------------------------------------|-----------|--------------------------------------|--------|--------------------------------------|
| America       | Fax: +1 919 878 3751                 |           | Fax: +44 (0)1865 852 739             |        | Fax: +49 (0) 89 8090 95 50           |
|               | Email: antibody_sales_us@bio-rad.com |           | Email: antibody_sales_uk@bio-rad.com |        | Email: antibody_sales_de@bio-rad.com |

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M421949:230815'

. . . . . .

#### Printed on 15 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint