Datasheet: MCA90T BATCH NUMBER 159864

Description:	MOUSE ANTI HUMAN CD90	
Specificity:	CD90	
Other names:	THY1	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	F15-42-1	
lsotype:	lgG1	
Quantity:	20 µg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>rad-antibodies.com/protocols.

	rad-antipodies.com/proto	COIS.						
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry	-			1/10 - 1/50			
	Immunohistology - Frozen (1)	-						
	Immunohistology - Paraffin							
	Immunoprecipitation	-						
	Immunofluorescence	-						
	Where this antibody has	not been	tested fo	r use in a particular teo	chnique this does not			
	necessarily exclude its us	necessarily exclude its use in such procedures. The suggested working dilution is given						
	as a guide only. It is recommended that the user titrates the antibody f system using appropriate negative/positive controls. (1)The epitope recognised by this antibody is reported to be sens formaldehyde fixation and tissue processing. Bio-Rad recommen acetone fixation for frozen sections.				sensitive to			
Target Species	Human							
Species Cross	Reacts with: Cynomolgus	s monkey	,					
Reactivity	N.B. Antibody reactivity a reactivity is derived from		-	• •	•			

Product Form	Purified IgG - liquid		
Preparation	Purified IgG prepared by ion exchange chromatography		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
Carrier Free	Yes		
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml		
Immunogen	Purified human brain Thy-1.		
External Database Links	UniProt: P04216 Related reagents Entrez Gene: 7070 THY1 Related reagents		
RRID	AB_871985		
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.		
Specificity	 Mouse anti Human CD90 antibody, clone F15-42-1 recognizes the human CD90 cell surface antigen, a ~25 kDa glycoprotein homologous to rat Thy1. The antigen is expressed by a subset of CD34+ve cells in the bone marrow and by prothymocytes within the thymus. CD90 is also expressed extensively within the brain. Mouse anti Human CD90 antibody, clone F15-42-1 is routinely tested in flow cytometry on the MOLT4 cell line. 		
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.		
Histology Positive Control Tissue	Human Brain, thymus		
References	 McKenzie, J.L. & Fabre, J.W. (1981) Human thy-1: unusual localization and possible functional significance in lymphoid tissues. <u>J Immunol. 126 (3): 843-50.</u> Daar, A.S. & Fabre, J.W. (1981) Demonstration with monoclonal antibodies of an 		

Cytochem. 52 (11): 1495-501.

5. Hagood, J.S. *et al.* (2005) Loss of fibroblast Thy-1 expression correlates with lung fibrogenesis. <u>Am J Pathol. 167 (2): 365-79.</u>

6. Diaz-Romero, J. *et al.* (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect bona fide dedifferentiation rather than amplification of progenitor cells. <u>J Cell Physiol. 214: 75-83.</u>

7. Cox, G. *et al.* (2011) The use of the reamer-irrigator-aspirator to harvest mesenchymal stem cells. J Bone Joint Surg Br. 93: 517-24.

8. Cizeau, J. *et al.* (2011) Fusogenics: a recombinant immunotoxin-based screening platform to select internalizing tumor-specific antibody fragments. <u>J Biomol Screen. 16:</u> <u>90-100.</u>

9. Gieseke, F. *et al.* (2010) Human multipotent mesenchymal stromal cells use galectin-1 to inhibit immune effector cells. <u>Blood. 116: 3770-9.</u>

10. Hauser, P.V. *et al.* (2010) Stem cells derived from human amniotic fluid contribute to acute kidney injury recovery. <u>Am J Pathol. 177: 2011-21.</u>

11. Holzwarth, C. *et al.* (2010) Low physiologic oxygen tensions reduce proliferation and differentiation of human multipotent mesenchymal stromal cells. BMC Cell Biol. 11:11

12. Karlsen, T.A. *et al.* (2010) Human primary articular chondrocytes, chondroblasts-like cells, and dedifferentiated chondrocytes: differences in gene, microRNA, and protein expression and phenotype. <u>Tissue Eng Part C Methods. 17: 219-27.</u>

13. Manochantr, S. *et al.* (2010) Isolation, characterization and neural differentiation potential of amnion derived mesenchymal stem cells. <u>J Med Assoc Thai. 93 Suppl 7:</u> <u>S183-91.</u>

14. Meng, J. *et al* (2011) Contribution of human muscle-derived cells to skeletal muscle regeneration in dystrophic host mice. <u>PLoS One. 6: e17454.</u>

15. Pessina, A. *et al.* (2010) CD45+/CD133+ positive cells expanded from umbilical cord blood expressing PDX-1 and markers of pluripotency. <u>Cell Biol Int. 34: 783-90.</u>

16. Tome, M. *et al.* (2007) Calponin is expressed by subpopulations of connective tissue cells but not olfactory ensheathing cells in the neonatal olfactory mucosa. <u>BMC Neurosci.</u> <u>8: 74.</u>

17. Yin, S. *et al.* (2010) Chondrogenic transdifferentiation of human dermal fibroblasts stimulated with cartilage-derived morphogenetic protein 1. <u>Tissue Eng Part A. 16:</u> <u>1633-43.</u>

18. Shafaei, H. *et al.* (2011) Effects of human placental serum on proliferation and morphology of human adipose tissue-derived stem cells. <u>Bone Marrow Transplant. 46:</u> <u>1464-71.</u>

19. Escobar, C.H. & Chaparro, O. (2016) Xeno-Free Extraction, Culture, and Cryopreservation of Human Adipose-Derived Mesenchymal Stem Cells. <u>Stem Cells Transl</u> <u>Med. 5 (3): 358-65.</u>

20. Shinoda, K. *et al.* (2016) Thy1+IL-7+ lymphatic endothelial cells in iBALT provide a survival niche for memory T-helper cells in allergic airway inflammation. <u>Proc Natl Acad</u> <u>Sci U S A. May 2. pii: 201512600. [Epub ahead of print]</u>

21. Kamprom, W. *et al.* (2016) Endothelial Progenitor Cell Migration-Enhancing Factors in the Secretome of Placental-Derived Mesenchymal Stem Cells. <u>Stem Cells Int. 2016:</u> <u>2514326.</u>

22. Vaquero, J. *et al.* (2016) An approach to personalized cell therapy in chronic complete paraplegia: The Puerta de Hierro phase I/II clinical trial. <u>Cytotherapy. 18 (8): 1025-36.</u>

	 Zhang, X. <i>et al.</i> (2017) Regeneration of hyaline-like cartilage in situ with SOX9 stimulation of bone marrow-derived mesenchymal stem cells. PLoS One. 12 (6): e0180138. GarikipatiV, N.S. <i>et al.</i> (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. PLoS One. 13 (2): e0192244. Chaturvedi, C.P. <i>et al.</i> (2018) Altered Expression of Hematopoiesis Regulatory Molecules in Lipopolysaccharide-Induced Bone Marrow Mesenchymal Stem Cells of Patients with Aplastic Anemia. Stem Cells Int. 2018: 6901761. Noda, S. <i>et al.</i> (2019) Effect of cell culture density on dental pulp-derived mesenchymal stem cells in Advanced Knee Osteoarthritis and Following Knee Joint Distraction. Front Bioeng Biotechnol. 8: 579751. Supokawej, A. <i>et al.</i> (2013) Cardiogenic and myogenic gene expression in mesenchymal stem cells after 5-azacytidine treatment. Turk J Haematol. 30 (2): 115-21. Paiboon, N. <i>et al.</i> (2019) MIF/CD74 axis participates in inflammatory activation of Schwann cells following sciatic nerve injury. J Mol Histol. 50 (4): 355-67. Fujii-Tezuka, R. <i>et al.</i> (2021) Umbilical artery tissue contains p75 neurotrophin receptor-positive pericyte-like cells that possess neurosphere formation capacity and neurogenic differentiation of pulse and rule capacity and neurogenic differentiation potential. <u>Regen Ther. 16: 1-11.</u> Orikasa, S. <i>et al.</i> (2022) Hypoxia-inducible factor 1α induces osteo/odontoblast differentiation of human dental pulp stem cells via Wnt/β-catenin transcriptional cofactor BCL9. Sci Rep. 12 (1): 682.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in
	frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA90T 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)RPEGoat Anti Mouse IgG IgA IgM (STAR87...)HRP

Goat Anti Mouse IgG (STAR76)	RPE
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>
Rabbit Anti Mouse IgG (STAR13)	HRP
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>
Goat Anti Mouse IgG (STAR77)	HRP
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,
	<u>DyLight®650</u> , <u>DyLight®680</u> , <u>DyLight®800</u> ,
	FITC, HRP
Recommended Negative Controls	

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-	rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M381517:210512'

Printed on 23 Jan 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint