

Datasheet: MCA90T

Description:	MOUSE ANTI HUMAN CD90
Specificity:	CD90
Other names:	THY1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	F15-42-1
Isotype:	IgG1
Quantity:	20 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
Immunoprecipitation	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. The suggested working dilution is given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species

Human

Species Cross Reactivity

Reacts with: Cynomolgus monkey

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

Preparation	Purified IgG prepared by ion exchange chromatography
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified human brain Thy-1.
External Database Links	<p>UniProt: P04216 Related reagents</p> <p>Entrez Gene: 7070 THY1 Related reagents</p>
RRID	AB_871985
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.
Specificity	<p>Mouse anti Human CD90 antibody, clone F15-42-1 recognizes the human CD90 cell surface antigen, a ~25 kDa glycoprotein homologous to rat Thy1. The antigen is expressed by a subset of CD34+ve cells in the bone marrow and by prothymocytes within the thymus. CD90 is also expressed extensively within the brain.</p> <p>Mouse anti Human CD90 antibody, clone F15-42-1 is routinely tested in flow cytometry on the MOLT4 cell line.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Human Brain, thymus
References	<ol style="list-style-type: none"> McKenzie, J.L. & Fabre, J.W. (1981) Human thy-1: unusual localization and possible functional significance in lymphoid tissues. J Immunol. 126 (3): 843-50. Daar, A.S. & Fabre, J.W. (1981) Demonstration with monoclonal antibodies of an unusual mononuclear cell infiltrate and loss of normal epithelial membrane antigens in human breast carcinomas. Lancet. 2 (8244): 434-8. Paul, G. <i>et al.</i> (2012) The adult human brain harbors multipotent perivascular mesenchymal stem cells. PLoS One. 7: e35577. Fiegel, H.C. <i>et al.</i> (2004) Stem-like cells in human hepatoblastoma. J Histochem Cytochem. 52 (11): 1495-501. Hagood, J.S. <i>et al.</i> (2005) Loss of fibroblast Thy-1 expression correlates with lung

- fibrogenesis. [Am J Pathol. 167 \(2\): 365-79.](#)
6. Diaz-Romero, J. *et al.* (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect bona fide dedifferentiation rather than amplification of progenitor cells. [J Cell Physiol. 214: 75-83.](#)
 7. Cox, G. *et al.* (2011) The use of the reamer-irrigator-aspirator to harvest mesenchymal stem cells. [J Bone Joint Surg Br. 93: 517-24.](#)
 8. Cizeau, J. *et al.* (2011) Fusogenics: a recombinant immunotoxin-based screening platform to select internalizing tumor-specific antibody fragments. [J Biomol Screen. 16: 90-100.](#)
 9. Gieseke, F. *et al.* (2010) Human multipotent mesenchymal stromal cells use galectin-1 to inhibit immune effector cells. [Blood. 116: 3770-9.](#)
 10. Hauser, P.V. *et al.* (2010) Stem cells derived from human amniotic fluid contribute to acute kidney injury recovery. [Am J Pathol. 177: 2011-21.](#)
 11. Holzwarth, C. *et al.* (2010) Low physiologic oxygen tensions reduce proliferation and differentiation of human multipotent mesenchymal stromal cells. [BMC Cell Biol. 11:11](#)
 12. Karlsen, T.A. *et al.* (2010) Human primary articular chondrocytes, chondroblasts-like cells, and dedifferentiated chondrocytes: differences in gene, microRNA, and protein expression and phenotype. [Tissue Eng Part C Methods. 17: 219-27.](#)
 13. Manochantr, S. *et al.* (2010) Isolation, characterization and neural differentiation potential of amnion derived mesenchymal stem cells. [J Med Assoc Thai. 93 Suppl 7: S183-91.](#)
 14. Meng, J. *et al.* (2011) Contribution of human muscle-derived cells to skeletal muscle regeneration in dystrophic host mice. [PLoS One. 6: e17454.](#)
 15. Pessina, A. *et al.* (2010) CD45+/CD133+ positive cells expanded from umbilical cord blood expressing PDX-1 and markers of pluripotency. [Cell Biol Int. 34: 783-90.](#)
 16. Tome, M. *et al.* (2007) Calponin is expressed by subpopulations of connective tissue cells but not olfactory ensheathing cells in the neonatal olfactory mucosa. [BMC Neurosci. 8: 74.](#)
 17. Yin, S. *et al.* (2010) Chondrogenic transdifferentiation of human dermal fibroblasts stimulated with cartilage-derived morphogenetic protein 1. [Tissue Eng Part A. 16: 1633-43.](#)
 18. Shafaei, H. *et al.* (2011) Effects of human placental serum on proliferation and morphology of human adipose tissue-derived stem cells. [Bone Marrow Transplant. 46: 1464-71.](#)
 19. Escobar, C.H. & Chaparro, O. (2016) Xeno-Free Extraction, Culture, and Cryopreservation of Human Adipose-Derived Mesenchymal Stem Cells. [Stem Cells Transl Med. 5 \(3\): 358-65.](#)
 20. Shinoda, K. *et al.* (2016) Thy1+IL-7+ lymphatic endothelial cells in iBALT provide a survival niche for memory T-helper cells in allergic airway inflammation. [Proc Natl Acad Sci U S A. May 2. pii: 201512600. \[Epub ahead of print\]](#)
 21. Kamprom, W. *et al.* (2016) Endothelial Progenitor Cell Migration-Enhancing Factors in the Secretome of Placental-Derived Mesenchymal Stem Cells. [Stem Cells Int. 2016: 2514326.](#)
 22. Vaquero, J. *et al.* (2016) An approach to personalized cell therapy in chronic complete paraplegia: The Puerta de Hierro phase I/II clinical trial. [Cytotherapy. 18 \(8\): 1025-36.](#)
 23. Zhang, X. *et al.* (2017) Regeneration of hyaline-like cartilage in situ with SOX9 stimulation of bone marrow-derived mesenchymal stem cells. [PLoS One. 12 \(6\):](#)

[e0180138](#).

24. GarikipatiV, N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. [PLoS One. 13 \(2\): e0192244](#).
25. Chaturvedi, C.P. *et al.* (2018) Altered Expression of Hematopoiesis Regulatory Molecules in Lipopolysaccharide-Induced Bone Marrow Mesenchymal Stem Cells of Patients with Aplastic Anemia. [Stem Cells Int. 2018: 6901761](#).
26. Noda, S. *et al.* (2019) Effect of cell culture density on dental pulp-derived mesenchymal stem cells with reference to osteogenic differentiation. [Sci Rep. 9 \(1\): 5430](#).
27. Sanjurjo-Rodriguez, C. *et al.* (2020) Gene Expression Signatures of Synovial Fluid Multipotent Stromal Cells in Advanced Knee Osteoarthritis and Following Knee Joint Distraction. [Front Bioeng Biotechnol. 8: 579751](#).
28. Supokawej, A. *et al.* (2013) Cardiogenic and myogenic gene expression in mesenchymal stem cells after 5-azacytidine treatment. [Turk J Haematol. 30 \(2\): 115-21](#).
29. Paiboon, N. *et al.* (2019) Gestational Tissue-Derived Human Mesenchymal Stem Cells Use Distinct Combinations of Bioactive Molecules to Suppress the Proliferation of Human Hepatoblastoma and Colorectal Cancer Cells. [Stem Cells Int. 2019: 9748795](#).
30. Song, H. *et al.* (2019) MIF/CD74 axis participates in inflammatory activation of Schwann cells following sciatic nerve injury. [J Mol Histol. 50 \(4\): 355-67](#).

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR8...)	DyLight@800
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@550 , DyLight@650 , DyLight@680 , DyLight@800 , FITC , HRP
Goat Anti Mouse IgG (STAR70...)	FITC

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M381517:210512'

Printed on 13 Oct 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)