Datasheet: MCA914 BATCH NUMBER 162716

Description:	MOUSE ANTI HUMAN CD55		
Specificity:	CD55		
Other names:	DAF		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	BRIC216		
Isotype:	lgG1		
Quantity:	0.2 mg		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

	rau-antipoules.com/proto	<u>cois</u> .				
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry					
	Immunohistology - Frozen			•		
	Immunohistology - Paraffin					
	ELISA			•		
	Immunoprecipitation					
	Western Blotting (1)					
	Functional Assays (2)					
	Where this antibody has not been tested for use in a particular technique this does not					
	necessarily exclude its use in such procedures. Suggested working dilutions are given					
	guide only. It is recommended that the user titrates the antibody for use in their own				or use in their own	
	system using appropriate negative/positive controls. (1) Non-reducing conditions required.					
	(2) This product contains sodium azide, removal by dialysis is recommended prio				s recommended prior	
	to use in functional ass	ays.				
Target Species	Human					
Product Form	Purified IgG - liquid					

 Preparation
 Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution	TRIS buffered saline.
Preservative Stabilisers	<0.1% Sodium Azide (NaN₃) ≤100mM Glycine
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Human fibroblast cell line.
External Database Links	UniProt: <u>P08174</u> <u>Related reagents</u> Entrez Gene: <u>1604</u> CD55 <u>Related reagents</u>
Synonyms	CR, DAF
RRID	AB_321792
Specificity	Mouse anti Human CD55 antibody, clone BRIC216 recognizes the CD55 antigen, a ~70 kDa glycoprotein also known as Decay Accelerating Factor (DAF). CD55 is distributed on erythrocytes and other circulating blood cells and also on cells in non-haemopoietic tissue particularly epithelium and endothelium. CD55 is also expressed at the foetal-maternal interfaces in placenta.CD55 has reduced expression on individuals with paroxysmal nocturnal haemoglobinuria. Mouse anti Human CD55 antibody, clone BRIC216 has a functional binding affinity to erythrocytes of 8.7 x 10^7 M ⁻¹ . The antigen is pronase and trypsin resistant and chymotrypsin sensitive. Mouse anti Human CD55 antibody, clone BRIC216 recognizes the consensus region 3 of the DAF molecule, which contains the functional site, and the antibody blocks the function of DAF.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul
References	 Fodor, W.L. <i>et al.</i> (1995) A novel bifunctional chimeric complement inhibitor that regulates C3 convertase and formation of the membrane attack complex. J Immunol. 155 (9): 4135-8. Wiesner, J. <i>et al.</i> (1997) Host cell factor CD59 restricts complement lysis of Plasmodium falciparum-infected erythrocytes. Eur J Immunol. 27 (10): 2708-13. Triantafilou, M. <i>et al.</i> (2000) A 70 kDa MHC class I associated protein (MAP-70) identified as a receptor molecule for Coxsackievirus A9 cell attachment. Hum Immunol. 61 (9): 867-78. Tieng, V. <i>et al.</i> (2002) Binding of Escherichia coli adhesin AfaE to CD55 triggers cell-surface expression of the MHC class I-related molecule MICA. Proc Natl Acad Sci U S A. 99: 2977-82. Loberg, R.D. <i>et al.</i> (2006) Inhibition of decay-accelerating factor (CD55) attenuates prostate cancer growth and survival <i>in vivo.</i> Neoplasia. 8: 69-78. Wu, G. <i>et al.</i> (2007) Coagulation cascade activation triggers early failure of pig hearts

	 expressing human complement regulatory genes. Xenotransplantation. 14 (1): 34-47. 7. Kim, M.S. & Racaniello, V.R. (2007) Enterovirus 70 receptor utilization is controlled by capsid residues that also regulate host range and cytopathogenicity. J Virol. 81 (16): 8648-55. 8. Ellison, B.S. <i>et al.</i> (2007) Complement susceptibility in glutamine deprived breast cancer cells. Cell Div. 2: 20. 9. Liszewski, M.K. <i>et al.</i> (2007) Modeling how CD46 deficiency predisposes to atypical hemolytic uremic syndrome. Mol Immunol. 44: 1559-68. 10. Koch, N. <i>et al.</i> (2009) IL-10 protects monocytes and macrophages from complement-mediated lysis. J Leukoc Biol. 86 (1): 155-66. 11. Tu, C.F. <i>et al.</i> (2010) The <i>in vitro</i> protection of human decay accelerating factor and hDAF/heme oxygenase-1 transgenes in porcine aortic endothelial cells against sera of Formosan macaques. Transplant Proc. 42 (6): 2138-41. 12. Pahwa, R. <i>et al.</i> (2018) Antibody Inhibition of Properdin Prevents Complement-Mediated Intravascular and Extravascular Hemolysis. J Immunol. 201 (3): 1021-9. 14. Noda, G.S. <i>et al.</i> (2020) Specificities and isotypes of erythrocytes autoantibodies in patients with warm autoimmune hemolytic anemia Rev Cubana Hematol Inmunol Hemoter 36(4): e1283
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10511 available at: https://www.bio-rad-antibodies.com/SDS/MCA914 10511
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77)	<u>HRP</u>		
Rabbit Anti Mouse IgG (STAR12)	RPE		
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>		
Goat Anti Mouse IgG IgA IgM (STAR87) Alk. Phos., HRP			
Goat Anti Mouse IgG (STAR76)	<u>RPE</u>		
Goat Anti Mouse IgG (Fc) (STAR120)	<u>FITC, HRP</u>		
Rabbit Anti Mouse IgG (STAR13)	<u>HRP</u>		
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>		

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

Alk. Phos., DyLight®488, DyLight®550, DyLight®650, DyLight®680, DyLight®800, FITC, HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide
America	Fax: +1 919 878 3751	
	Email: antibody_sales_us@bio-rad	.com

 Tel: +44 (0)1865 852 700
 Europe

 Fax: +44 (0)1865 852 739
 Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M390514:210915'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint