

## Datasheet: MCA928APC BATCH NUMBER 155309

Description:	MOUSE IgG1 NEGATIVE CONTROL:APC
Specificity:	MOUSE IgG1 NEGATIVE CONTROL
Format:	APC
Product Type:	Negative/Isotype Control
Isotype:	lgG1
Quantity:	100 TESTS

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes No	Not Determined	Suggested Dilution			
	Flow Cytometry	•		*			
	necessarily exclude it	ts use in such proc commended that th	e user titrates the antibo	king dilutions are given as			
Target Species	Negative Control						
Product Form	Purified IgG conjugated to APC- lyophilised						
Reconstitution	Reconstitute with 1.0 ml distilled water						
Max Ex/Em	Fluorophore	Excitation Max (n	m) Emission Max (nm)				
	APC	650	661	_			
Preparation	Purified IgG prepared by affinity chromatography on Protein A						
Buffer Solution	Phosphate buffered saline						
Preservative	0.09% Sodium Azide						
Stabilisers	1% Bovine Serum	Albumin					
	5% Sucrose						
RRID	AB_322309						

SpecificityMouse IgG1 negative control is negative by flow cytometry on all human cells and cell<br/>lines tested. Further tests have also shown that this reagent is also suitable for use as a<br/>negative control with bovine (Maslanka *et al*, 2012), ovine, porcine (Kapetanovic *et al*,<br/>2012), equine (Jacks *et al*, 2007), canine (Maiolini *et al*, 2012), lapine (Pakandl *et al*,<br/>2008) and guinea-pig tissues.

## This reagent recognizes a rat cell surface marker, and therefore cannot be used as a negative control in this species.

Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells or cells or 100ul whole blood
References	1. Kupatt, C. <i>et al.</i> (2000) c7E3Fab reduces postischemic leukocyte-thrombocyte interaction mediated by fibrinogen. Implications for myocardial reperfusion injury. <u>Arterioscler Thromb Vasc Biol. 20 (10): 2226-32.</u>
	2. Dalli, J. <i>et al.</i> (2008) Annexin 1 mediates the rapid anti-inflammatory effects of
	neutrophil-derived microparticles. <u>Blood. 112 (6): 2512-9.</u>
	3. Barratt-Due, A. <i>et al.</i> (2011) Ornithodoros moubata Complement Inhibitor Is an Equally
	Effective C5 Inhibitor in Pigs and Humans. <u>J Immunol. 187: 4913-9.</u>
	4. Kapetanovic, R. et al. (2012) Pig bone marrow-derived macrophages resemble human
	macrophages in their response to bacterial lipopolysaccharide. J Immunol. 188: 3382-94.
	5. Maiolini, A. et al. (2012) Toll-like receptors 4 and 9 are responsible for the maintenance
	of the inflammatory reaction in canine steroid-responsive meningitis-arteritis, a large
	animal model for neutrophilic meningitis. J Neuroinflammation. 9: 226.
	6. Maślanka, T. <i>et al.</i> (2012) The presence of CD25 on bovine WC1+ gammadelta T cells
	is positively correlated with their production of IL-10 and TGF-beta, but not IFN-gamma.
	Pol J Vet Sci. 15 (1): 11-20.
	7. Pakandl, M. et al. (2008) Immune response to rabbit coccidiosis: a comparison betwee
	infections with Eimeria flavescens and E. intestinalis. Folia Parasitol (Praha). 55:1-6.
	8. Jacks, S. et al. (2007) Experimental infection of neonatal foals with Rhodococcus equi
	triggers adult-like gamma interferon induction. Clin Vaccine Immunol.14:669-77
	9. Kamble, N.M. et al. (2016) Interaction of a live attenuated Salmonella gallinarum
	vaccine candidate with chicken bone marrow-derived dendritic cells. <u>Avian Pathol. Jan</u>
	26:1-24. [Epub ahead of print]
	10. Brace, P.T. <i>et al.</i> (2017) <i>Mycobacterium tuberculosis</i> subverts negative regulatory
	pathways in human macrophages to drive immunopathology. <u>PLoS Pathog. 13 (6):</u>
	e1006367.
	11. Topoluk, N. <i>et al.</i> (2017) Amniotic Mesenchymal Stromal Cells Exhibit Preferential
	Osteogenic and Chondrogenic Differentiation and Enhanced Matrix Production Compared
	With Adipose Mesenchymal Stromal Cells. <u>Am J Sports Med. 363546517706138.</u>
	12. Iwaszko-Simonik, A. <i>et al.</i> (2015) Expression of surface platelet receptors (CD62P an
	CD41/61) in horses with recurrent airway obstruction (RAO). <u>Vet Immunol Immunopathol</u> .
	<u>164 (1-2): 87-92.</u> 13. Arzi, B. <i>et al.</i> (2017) Therapeutic Efficacy of Fresh, Allogeneic Mesenchymal Stem
	Cells for Severe Refractory Feline Chronic Gingivostomatitis. <u>Stem Cells Transl Med. 6</u>
	(8): 1710-22.
	14. Taechangam, N. <i>et al.</i> (2021) Feline adipose-derived mesenchymal stem cells induce
	effector phenotype and enhance cytolytic function of CD8+ T cells. <u>Stem Cell Res Ther.</u>

		<u>12 (1): 49</u>	95.				
Storage		Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.					
Guarante	e	12 months from date of despatch					
Health And Safety Information		Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA928APC 20487					
Regulato	ry	For resea	arch purposes	s only			
North & South America	Tel: +1 800 265 73 Fax: +1 919 878 3 Email: antibody_s	751	Worldwide rad.com	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bi	Europe o-rad.com	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com	
To find a b	atch/lot specifio	c datashee	t for this produ	ict, please use our online 'M375697:210104'	search tool at: b	vio-rad-antibodies.com/datasheets	
				Printed on 29 Feb 2024			

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint