

## Datasheet: MCA928B

<b>Description:</b>	MOUSE IgG1 NEGATIVE CONTROL:Biotin
<b>Specificity:</b>	MOUSE IgG1 NEGATIVE CONTROL
<b>Format:</b>	Biotin
<b>Product Type:</b>	Negative/Isotype Control
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS

## Product Details

**RRID** AB\_322261

**Applications** This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. \*It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test reagents.

**Target Species** Negative Control

**Product Form** Purified IgG conjugated to Biotin - liquid

**Preparation** Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

**Buffer Solution** Phosphate buffered saline

**Preservative** 0.09% Sodium Azide  
**Stabilisers** 1% Bovine Serum Albumin

**Approx. Protein Concentrations** IgG concentration 0.1 mg/ml

**Specificity** **Mouse IgG1 negative control** is negative by flow cytometry on all human cells and cell lines tested. Further tests have also shown that this reagent is also suitable for use as a negative control with bovine ([Maslanka et al. 2012](#)), ovine, porcine ([Kapetanovic et al. 2012](#)), equine ([Jacks et al. 2007](#)), canine ([Maiolini et al. 2012](#)), lapine ([Pakandl et al. 2008](#)) and guinea-pig tissues.

***This reagent recognizes a rat cell surface marker, and therefore cannot be used as a negative control in this species.***

**Flow Cytometry** Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells or 100ul whole blood.

## References

1. Kupatt, C. *et al.* (2000) c7E3Fab reduces postischemic leukocyte-thrombocyte interaction mediated by fibrinogen. Implications for myocardial reperfusion injury. [Arterioscler Thromb Vasc Biol. 20 \(10\): 2226-32.](#)
2. Dalli, J. *et al.* (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. [Blood. 112 \(6\): 2512-9.](#)
3. Barratt-Due, A. *et al.* (2011) Ornithodoros moubata Complement Inhibitor Is an Equally Effective C5 Inhibitor in Pigs and Humans. [J Immunol. 187: 4913-9.](#)
4. Kapetanovic, R. *et al.* (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. [J Immunol. 188: 3382-94.](#)
5. Maiolini, A. *et al.* (2012) Toll-like receptors 4 and 9 are responsible for the maintenance of the inflammatory reaction in canine steroid-responsive meningitis-arteritis, a large animal model for neutrophilic meningitis. [J Neuroinflammation. 9: 226.](#)
6. Maślanka, T. *et al.* (2012) The presence of CD25 on bovine WC1+  $\gamma\delta$  T cells is positively correlated with their production of IL-10 and TGF- $\beta$ , but not IFN- $\gamma$  [Polish Journal of Veterinary Sciences.15: 11–20.](#)
7. Pakandl, M. *et al.* (2008) Immune response to rabbit coccidiosis: a comparison between infections with Eimeria flavescens and E. intestinalis. [Folia Parasitol \(Praha\). 55:1-6.](#)
8. Jacks, S. *et al.* (2007) Experimental infection of neonatal foals with Rhodococcus equi triggers adult-like gamma interferon induction. [Clin Vaccine Immunol.14:669-77](#)
9. Kamble, N.M. *et al.* (2016) Interaction of a live attenuated *Salmonella gallinarum* vaccine candidate with chicken bone marrow-derived dendritic cells. [Avian Pathol. Jan 26:1-24. \[Epub ahead of print\]](#)
10. Brace, P.T. *et al.* (2017) *Mycobacterium tuberculosis* subverts negative regulatory pathways in human macrophages to drive immunopathology. [PLoS Pathog. 13 \(6\): e1006367.](#)
11. Topoluk, N. *et al.* (2017) Amniotic Mesenchymal Stromal Cells Exhibit Preferential Osteogenic and Chondrogenic Differentiation and Enhanced Matrix Production Compared With Adipose Mesenchymal Stromal Cells. [Am J Sports Med. 363546517706138.](#)
12. Iwaszko-Simonik, A. *et al.* (2015) Expression of surface platelet receptors (CD62P and CD41/61) in horses with recurrent airway obstruction (RAO). [Vet Immunol Immunopathol. 164 \(1-2\): 87-92.](#)
13. Arzi, B. *et al.* (2017) Therapeutic Efficacy of Fresh, Allogeneic Mesenchymal Stem Cells for Severe Refractory Feline Chronic Gingivostomatitis. [Stem Cells Transl Med. 6 \(8\): 1710-22.](#)

---

### Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

### Guarantee

12 months from date of reconstitution.

---

### Health And Safety Information

Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

---

### Regulatory

For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Biotin \(MCA1209B\)](#)

**North & South America** Tel: +1 800 265 7376  
Fax: +1 919 878 3751  
Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide** Tel: +44 (0)1865 852 700  
Fax: +44 (0)1865 852 739  
Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)  
'M335670:181204'

**Europe** Tel: +49 (0) 89 8090 95 21  
Fax: +49 (0) 89 8090 95 50  
Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

**Printed on 11 Oct 2019**

---

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)