

Datasheet: MCA928EL

BATCH NUMBER 166171

Description:	MOUSE IgG1 NEGATIVE CONTROL:Low Endotoxin
Specificity:	MOUSE IgG1 NEGATIVE CONTROL
Format:	Low Endotoxin
Product Type:	Negative/Isotype Control
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			*
Immunohistology - Frozen	■			
Immunohistology - Paraffin			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. * It is recommended that the user titrates the antibody for use in their own system to a concentration equivalent to their test reagents.

Target Species	Negative Control
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Carrier Free	Yes
Endotoxin Level	<0.01 EU/ug
Approx. Protein	IgG concentration 1.0 mg/ml

Concentrations

RRID AB_324168

Specificity **Mouse IgG1 negative control** is negative by flow cytometry on all human cells and cell lines tested. Further tests have also shown that this reagent is also suitable for use as a negative control with bovine (Maslanka *et al*, 2012), ovine, porcine (Kapetanovic *et al*, 2012), equine (Jacks *et al*, 2007), canine (Maiolini *et al*, 2012), lapine (Pakandl *et al*, 2008) and guinea-pig tissues.

This reagent recognizes a rat cell surface marker, and therefore cannot be used as a negative control in this species.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

- References**
1. Kupatt, C. *et al*. (2000) c7E3Fab reduces postischemic leukocyte-thrombocyte interaction mediated by fibrinogen. Implications for myocardial reperfusion injury. [Arterioscler Thromb Vasc Biol. 20 \(10\): 2226-32.](#)
 2. Jacks, S. *et al*. (2007) Experimental infection of neonatal foals with *Rhodococcus equi* triggers adult-like gamma interferon induction. [Clin Vaccine Immunol. 14:669-77](#)
 3. Pakandl, M. *et al*. (2008) Immune response to rabbit coccidiosis: a comparison between infections with *Eimeria flavescens* and *E. intestinalis*. [Folia Parasitol \(Praha\). 55:1-6.](#)
 4. Dalli, J. *et al*. (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. [Blood. 112 \(6\): 2512-9.](#)
 5. Barratt-Due, A. *et al*. (2011) *Ornithodoros moubata* Complement Inhibitor Is an Equally Effective C5 Inhibitor in Pigs and Humans. [J Immunol. 187: 4913-9.](#)
 6. Maślanka, T. *et al*. (2012) The presence of CD25 on bovine WC1+ gammadelta T cells is positively correlated with their production of IL-10 and TGF-beta, but not IFN-gamma. [Pol J Vet Sci. 15 \(1\): 11-20.](#)
 7. Maiolini, A. *et al*. (2012) Toll-like receptors 4 and 9 are responsible for the maintenance of the inflammatory reaction in canine steroid-responsive meningitis-arteritis, a large animal model for neutrophilic meningitis. [J Neuroinflammation. 9: 226.](#)
 8. Kapetanovic, R. *et al*. (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. [J Immunol. 188: 3382-94.](#)
 9. Kamble, N.M. *et al*. (2016) Interaction of a live attenuated *Salmonella Gallinarum* vaccine candidate with chicken bone marrow-derived dendritic cells. [Avian Pathol. 45 \(2\): 235-43.](#)
 10. Iwaszko-Simonik, A. *et al*. (2015) Expression of surface platelet receptors (CD62P and CD41/61) in horses with recurrent airway obstruction (RAO). [Vet Immunol Immunopathol. 164 \(1-2\): 87-92.](#)
 11. Brace, P.T. *et al*. (2017) *Mycobacterium tuberculosis* subverts negative regulatory pathways in human macrophages to drive immunopathology. [PLoS Pathog. 13 \(6\): e1006367.](#)
 12. Topoluk, N. *et al*. (2017) Amniotic Mesenchymal Stromal Cells Exhibit Preferential Osteogenic and Chondrogenic Differentiation and Enhanced Matrix Production Compared With Adipose Mesenchymal Stromal Cells. [Am J Sports Med. 45 \(11\): 2637-46.](#)
 13. Arzi, B. *et al*. (2017) Therapeutic Efficacy of Fresh, Allogeneic Mesenchymal Stem Cells for Severe Refractory Feline Chronic Gingivostomatitis. [Stem Cells Transl Med. 6](#)

(8): 1710-22.

14. Taechangam, N. *et al.* (2021) Feline adipose-derived mesenchymal stem cells induce effector phenotype and enhance cytolytic function of CD8+ T cells. [Stem Cell Res Ther. 12 \(1\): 495.](#)

15. do Prado Duzanski, A.*et al.* (2022) Cell-mediated immunity and expression of MHC class I and class II molecules in dogs naturally infected by canine transmissible venereal tumor: Is there complete spontaneous regression outside the experimental CTVT? [Research in Veterinary Science. 145: 193-204.](#)

16. Tolstova, T. *et al.* (2023) The effect of TLR3 priming conditions on MSC immunosuppressive properties. [Stem Cell Res Ther. 14 \(1\): 344.](#)

17. Geng, Y. *et al.* (2018) Dietary vitamin D(3) supplementation protects laying hens against lipopolysaccharide-induced immunological stress. [Nutr Metab \(Lond\). 15: 58.](#)

18. Dan-Jumbo, S.O. *et al.* (2024) Derivation and long-term maintenance of porcine skeletal muscle progenitor cells. [Sci Rep. 14 \(1\): 9370.](#)

Storage

Store at -20°C only.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10162 available at: <https://www.bio-rad-antibodies.com/SDS/MCA928EL>
10162

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Low Endotoxin \(MCA1209EL\)](#)

North & South America

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M369175:200529'

Printed on 09 May 2024