



Datasheet: MCA928PB

BATCH NUMBER 166047

Description:	MOUSE IgG1 NEGATIVE CONTROL:Pacific Blue®
Specificity:	MOUSE IgG1 NEGATIVE CONTROL
Format:	Pacific Blue®
Product Type:	Negative/Isotype Control
Isotype:	IgG1
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures.

*This antibody should be used at the same concentration as the test antibody.

Target Species	Negative Control						
Product Form	Purified IgG conjugated to Pacific Blue® - liquid						
Max Ex/Em	<table><thead><tr><th>Fluorophore</th><th>Excitation Max (nm)</th><th>Emission Max (nm)</th></tr></thead><tbody><tr><td>Pacific Blue®</td><td>410</td><td>455</td></tr></tbody></table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	Pacific Blue®	410	455
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
Pacific Blue®	410	455					
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin						
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml						
RRID	AB_567338						

Specificity

Mouse IgG1 negative control is negative by flow cytometry on all human cells and cell lines tested. Further tests have also shown that this reagent is also suitable for use as a negative control with bovine (Maslanka *et al*, 2012), ovine, porcine ([Kapetanovic *et al*, 2012](#)), equine ([Jacks *et al*, 2007](#)), canine ([Maiolini *et al*, 2012](#)), lapine ([Pakandl *et al*, 2008](#)) and guinea-pig tissues.

This reagent recognizes a rat cell surface marker, and therefore cannot be used as a negative control in this species.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells or 100ul whole blood.

References

1. Kupatt, C. *et al*. (2000) c7E3Fab reduces postischemic leukocyte-thrombocyte interaction mediated by fibrinogen. Implications for myocardial reperfusion injury. [Arterioscler Thromb Vasc Biol. 20 \(10\): 2226-32.](#)
2. Jacks, S. *et al*. (2007) Experimental infection of neonatal foals with *Rhodococcus equi* triggers adult-like gamma interferon induction. [Clin Vaccine Immunol. 14:669-77](#)
3. Pakandl, M. *et al*. (2008) Immune response to rabbit coccidiosis: a comparison between infections with *Eimeria flavescens* and *E. intestinalis*. [Folia Parasitol \(Praha\). 55:1-6.](#)
4. Dalli, J. *et al*. (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. [Blood. 112 \(6\): 2512-9.](#)
5. Barratt-Due, A. *et al*. (2011) *Ornithodoros moubata* Complement Inhibitor Is an Equally Effective C5 Inhibitor in Pigs and Humans. [J Immunol. 187: 4913-9.](#)
6. Maślanka, T. *et al*. (2012) The presence of CD25 on bovine WC1+ gammadelta T cells is positively correlated with their production of IL-10 and TGF-beta, but not IFN-gamma. [Pol J Vet Sci. 15 \(1\): 11-20.](#)
7. Maiolini, A. *et al*. (2012) Toll-like receptors 4 and 9 are responsible for the maintenance of the inflammatory reaction in canine steroid-responsive meningitis-arteritis, a large animal model for neutrophilic meningitis. [J Neuroinflammation. 9: 226.](#)
8. Kapetanovic, R. *et al*. (2012) Pig bone marrow-derived macrophages resemble human macrophages in their response to bacterial lipopolysaccharide. [J Immunol. 188: 3382-94.](#)
9. Kamble, N.M. *et al*. (2016) Interaction of a live attenuated *Salmonella Gallinarum* vaccine candidate with chicken bone marrow-derived dendritic cells. [Avian Pathol. 45 \(2\): 235-43.](#)
10. Iwaszko-Simonik, A. *et al*. (2015) Expression of surface platelet receptors (CD62P and CD41/61) in horses with recurrent airway obstruction (RAO). [Vet Immunol Immunopathol. 164 \(1-2\): 87-92.](#)
11. Brace, P.T. *et al*. (2017) *Mycobacterium tuberculosis* subverts negative regulatory pathways in human macrophages to drive immunopathology. [PLoS Pathog. 13 \(6\): e1006367.](#)
12. Topoluk, N. *et al*. (2017) Amniotic Mesenchymal Stromal Cells Exhibit Preferential Osteogenic and Chondrogenic Differentiation and Enhanced Matrix Production Compared With Adipose Mesenchymal Stromal Cells. [Am J Sports Med. 45 \(11\): 2637-46.](#)
13. Arzi, B. *et al*. (2017) Therapeutic Efficacy of Fresh, Allogeneic Mesenchymal Stem Cells for Severe Refractory Feline Chronic Gingivostomatitis. [Stem Cells Transl Med. 6 \(8\): 1710-22.](#)
14. Taechangam, N. *et al*. (2021) Feline adipose-derived mesenchymal stem cells induce effector phenotype and enhance cytolytic function of CD8+ T cells. [Stem Cell Res Ther.](#)

[12 \(1\): 495.](#)

15. do Prado Duzanski, A. *et al.* (2022) Cell-mediated immunity and expression of MHC class I and class II molecules in dogs naturally infected by canine transmissible venereal tumor: Is there complete spontaneous regression outside the experimental CTVT?

[Research in Veterinary Science. 145: 193-204.](#)

16. Tolstova, T. *et al.* (2023) The effect of TLR3 priming conditions on MSC immunosuppressive properties. [Stem Cell Res Ther. 14 \(1\): 344.](#)

17. Geng, Y. *et al.* (2018) Dietary vitamin D(3) supplementation protects laying hens against lipopolysaccharide-induced immunological stress. [Nutr Metab \(Lond\). 15: 58.](#)

18. Dan-Jumbo, S.O. *et al.* (2024) Derivation and long-term maintenance of porcine skeletal muscle progenitor cells. [Sci Rep. 14 \(1\): 9370.](#)

19. Maciag, S. *et al.* (2022) Effects of freezing storage on the stability of maternal cellular and humoral immune components in porcine colostrum. [Vet Immunol Immunopathol. 254: 110520.](#)

20. Forner, R. *et al.* (2021) Distribution difference of colostrum-derived B and T cells subsets in gilts and sows. [PLoS One. 16 \(5\): e0249366.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee

12 months from date of despatch

Acknowledgements

The Pacific Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays, and are covered by pending and issued patents.

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA928PB>
10041

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Pacific Blue® \(MCA1209PB\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M384837:210513'

Printed on 28 Aug 2024

