

Datasheet: MCA929B

Description:	MOUSE IgG2a NEGATIVE CONTROL:Biotin
Specificity:	MOUSE IgG2a NEGATIVE CONTROL
Format:	Biotin
Product Type:	Negative/Isotype Control
Isotype:	lgG2a
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			*
Immunohistology - Frozen			•	
Immunohistology - Paraffin				
ELISA				
Western Blotting			•	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures.

*It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test reagents.

Target Species	Negative Control
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Activated rat T-helper cells.

	pleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse nyeloma cell line.
Co	louse IgG2a negative control antibody, clone OX-34 is suitable for use as a negative ontrol reagent for the measurement of non-specific binding of mouse monoclonal ntibodies of isotype IgG2a to human tissue.
	clone MRC OX-34 recognises a rat cell surface marker, and therefore cannot be used as negative control in this species.
	this product is routinely tested in flow cytometry on rat splenocytes to confirm antibody ctivity and on human whole blood to test for suitability as a negative control.
	est results have shown that MCA929 is also suitable for use as a negative control with ovine, ovine, porcine, equine, canine, lapine and guinea-pig tissues.
Т	his antibody may not be suitable for intracellular staining on some cell types.
Flow Cytometry U	lse 10ul of the suggested working dilution to label 10 ⁶ in 100ul.
tr 2 v 2 2 3	. Avigdor, A. et al. (2004) CD44 and hyaluronic acid cooperate with SDF-1 in the afficking of human CD34+ stem/progenitor cells to bone marrow. Blood. 103 (8): 2981-9. . Kamble, N.M. et al. (2016) Interaction of a live attenuated Salmonella gallinarum accine candidate with chicken bone marrow-derived dendritic cells. Avian Pathol. Jan 6:1-24. [Epub ahead of print] . Wattegedera, S.R. et al. (2017) Enhancing the toolbox to study IL-17A in cattle and heep. Vet Res. 48 (1): 20.
-2	This product is shipped at ambient temperature. It is recommended to aliquot and store at 20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for hort term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	void repeated freezing and thawing as this may denature the antibody. Storage in ost-free freezers is not recommended.
Guarantee 1	2 months from date of despatch
Information <u>h</u>	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA929B
Regulatory F	or research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

Printed on 25 Mar 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint