

Datasheet: MCA929EL

BATCH NUMBER 0714

Description:	MOUSE IgG2a NEGATIVE CONTROL:Low Endotoxin			
Specificity:	MOUSE IgG2a NEGATIVE CONTROL			
Format:	Low Endotoxin			
Product Type:	Negative/Isotype Control			
Isotype:	lgG2a			
Quantity:	0.5 mg			

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	•			*		
	Immunohistology - Frozen			-			
	Immunohistology - Paraffin			•			
	Where this antibody has not been tested for use in a particular technique this does not						
	necessarily exclude its us	necessarily exclude its use in such procedures.					
	* It is recommended that	the user of	dilutes the	antibody to a concer	ntration equivalent to		
	their test reagent.						
Target Species	Negative Control						
Product Form	Purified IgG - liquid						
Preparation	Purified IgG prepared by supernatant	affinity ch	nromatogra	aphy on Protein A fro	m tissue culture		
Buffer Solution	Phosphate buffered salin	е					
Preservative Stabilisers	None present						
Carrier Free	Yes						
Endotoxin Level	<0.01 EU/mg						
Approx. Protein	IgG concentration 1.0 mg	g/ml					

Concentrations

Immunogen	Activated rat T-helper cells.				
RRID	AB_567348				
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.				
Specificity	Mouse IgG2a negative control antibody, clone OX-34 is suitable for use as a negative control reagent for the measurement of non-specific binding of mouse monoclonal antibodies of isotype IgG2a to human tissue.				
	Clone MRC OX-34 recognises a rat cell surface marker, and therefore cannot be used as a negative control in this species.				
	This product is routinely tested in flow cytometry on rat splenocytes to confirm antibody activity and on human whole blood to test for suitability as a negative control.				
	Test results have shown that MCA929 is also suitable for use as a negative control with bovine, ovine, porcine, equine, canine, lapine and guinea-pig tissues.				
	This antibody may not be suitable for intracellular staining on some cell types.				
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.				
References	 Avigdor, A. <i>et al.</i> (2004) CD44 and hyaluronic acid cooperate with SDF-1 in the trafficking of human CD34+ stem/progenitor cells to bone marrow. <u>Blood. 103 (8): 2981-9.</u> Kamble, N.M. <i>et al.</i> (2016) Interaction of a live attenuated Salmonella Gallinarum vaccine candidate with chicken bone marrow-derived dendritic cells. <u>Avian Pathol. 45 (2): 235-43.</u> Wattegedera, S.R. <i>et al.</i> (2017) Enhancing the toolbox to study IL-17A in cattle and sheep. <u>Vet Res. 48 (1): 20.</u> Terpeluk, R.E. <i>et al.</i> (2024) Supplementation of Foals with a <i>Saccharomyces cerevisiae</i> Fermentation Product Alters the Early Response to Vaccination <u>Animals. 14 (6): 960.</u> Maciag, S. <i>et al.</i> (2022) Effects of freezing storage on the stability of maternal cellular and humoral immune components in porcine colostrum. <u>Vet Immunol Immunopathol. 254: 110520.</u> Forner, R. <i>et al.</i> (2021) Distribution difference of colostrum-derived B and T cells subsets in gilts and sows. <u>PLoS One. 16 (5): e0249366.</u> Rogato, F. <i>et al.</i> (2024) Leukemia cutis as a prominent clinical sign in a dog with acute myeloid leukemia. <u>Vet Clin Pathol. 53 (4): 448-57.</u> 				
Storage	Store at -20°C only.				
	This product should be stored undiluted.				
	Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing				

			in a precipitate we			
Guarante	e					
Information <u>ht</u>			l Safety Datas /ww.bio-rad-ar	ıt:		
Regulato	ry	For rese	earch purpose	s only		
orth & South	Tel: +1 800 265 Fax: +1 919 87		Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
	Email: antibody	_sales_us@bi	o-rad.com	Email: antibody_sales_uk@bic	-rad.com	Email: antibody_sales_de@bio-rad.com
To find a b	atch/lot spec	ific datashe	et for this produ	uct, please use our online 'M369191:200529'	search tool at: <mark>b</mark> i	o-rad-antibodies.com/datashee
				Printed on 27 Mar 2025		

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint