

Datasheet: MCA929F BATCH NUMBER 0515

Description:	MOUSE IgG2a NEGATIVE CONTROL:FITC
Specificity:	MOUSE IgG2a NEGATIVE CONTROL
Format:	FITC
Product Type:	Negative/Isotype Control
Isotype:	lgG2a
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				*

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures.

^{*} It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test reagent.

Target Species	Negative Control				
Product Form	Purified IgG conjugat	niocyanate Isomer 1 (FI	1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	FITC	490	525		
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant				
Buffer Solution	Phosphate buffered saline				
Preservative	0.09% Sodium Azide	•			
Stabilisers	1% Bovine Serum				
Approx. Protein Concentrations	IgG concentration 0.	1 mg/ml			
Immunogen	Activated rat T-helper cells.				

Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.
Specificity	Mouse IgG2a negative control antibody, clone OX-34 is suitable for use as a negative control reagent for the measurement of non-specific binding of mouse monoclonal antibodies of isotype IgG2a to human tissue.
	Clone MRC OX-34 recognises a rat cell surface marker, and therefore cannot be used as a negative control in this species.
	This product is routinely tested in flow cytometry on rat splenocytes to confirm antibody activity and on human whole blood to test for suitability as a negative control.
	Test results have shown that MCA929 is also suitable for use as a negative control with bovine, ovine, porcine, equine, canine, lapine and guinea-pig tissues.
	This antibody may not be suitable for intracellular staining on some cell types.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Avigdor, A. <i>et al.</i> (2004) CD44 and hyaluronic acid cooperate with SDF-1 in the trafficking of human CD34+ stem/progenitor cells to bone marrow. <u>Blood. 103 (8): 2981-9981-9981.</u> 2. Kamble, N.M. <i>et al.</i> (2016) Interaction of a live attenuated <i>Salmonella gallinarum</i> vaccine candidate with chicken bone marrow-derived dendritic cells. <u>Avian Pathol. Jan 26:1-24. [Epub ahead of print]</u> 3. Wattegedera, S.R. <i>et al.</i> (2017) Enhancing the toolbox to study IL-17A in cattle and sheep. <u>Vet Res. 48 (1): 20.</u>
Storage	Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted.
	Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
	Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA929F 10041

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Europe

'M302087:170109'

Email: antibody_sales_de@bio-rad.com

Printed on 25 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint