

## Datasheet: MCA929PE BATCH NUMBER 160695

Description:	MOUSE IgG2a NEGATIVE CONTROL:RPE
Specificity:	MOUSE IgG2a NEGATIVE CONTROL
Format:	RPE
Product Type:	Negative/Isotype Control
Isotype:	lgG2a
Quantity:	100 TESTS

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes I	No Not Determ	nined Suggested Dil	lution		
	Flow Cytometry	•		*			
	necessarily exclude its	ocedures. Suggeste the user dilutes the	cular technique this does d working dilutions are giv e antibody for use in their o	ven as			
Target Species	Negative Control						
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized						
Reconstitution	Reconstitute with 1.0						
Max Ex/Em	Fluorophore	Excitation Max	(nm) Emission Ma	x (nm)			
	RPE 488nm laser	496	578				
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative	0.09% Sodium Azide						
Stabilisers	1% Bovine Serum Albumin						
	5% Sucrose						

Immunogen	Activated rat T-helper cells.
RRID	AB_322272
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.
Specificity	<b>Mouse IgG2a negative control antibody, clone OX-34</b> is suitable for use as a negative control reagent for the measurement of non-specific binding of mouse monoclonal antibodies of isotype IgG2a to human tissue.
	Clone MRC OX-34 recognises a rat cell surface marker, and therefore cannot be used as a negative control in this species.
	This product is routinely tested in flow cytometry on rat splenocytes to confirm antibody activity and on human whole blood to test for suitability as a negative control.
	Test results have shown that MCA929 is also suitable for use as a negative control with bovine, ovine, porcine, equine, canine, lapine and guinea-pig tissues.
	This antibody may not be suitable for intracellular staining on some cell types.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
References	<ol> <li>Avigdor, A. <i>et al.</i> (2004) CD44 and hyaluronic acid cooperate with SDF-1 in the trafficking of human CD34+ stem/progenitor cells to bone marrow. <u>Blood. 103 (8): 2981-9.</u></li> <li>Kamble, N.M. <i>et al.</i> (2016) Interaction of a live attenuated <i>Salmonella gallinarum</i> vaccine candidate with chicken bone marrow-derived dendritic cells. <u>Avian Pathol. Jan 26:1-24. [Epub ahead of print]</u></li> <li>Wattegedera, S.R. <i>et al.</i> (2017) Enhancing the toolbox to study IL-17A in cattle and sheep. <u>Vet Res. 48 (1): 20.</u></li> </ol>
Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.
	DO NOT FREEZE.
	This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA929PE 20487
Regulatory	For research purposes only

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21			
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50	То		
	Email: antibody_sales_us@bio-r	ad.com	Email: antibody_sales_uk@bio-	rad.com	Email: antibody_sales_de@bio-ra	<sup>id.</sup> md a		
batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375703:210104'								

## Printed on 25 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint