

## Datasheet: MCA986AMO

**BATCH NUMBER 149338**

<b>Description:</b>	MOUSE ANTI HUMAN HLA B7:Amethyst Orange
<b>Specificity:</b>	HLA B7
<b>Format:</b>	Amethyst Orange
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	BB7.1
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

#### Target Species

Human

#### Species Cross Reactivity

Reacts with: Cynomolgus monkey

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG conjugated to Amethyst Orange - liquid

#### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
Amethyst Orange	405	540

#### Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

#### Buffer Solution

Phosphate buffered saline

<b>Preservative</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Stabilisers</b>	1% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml
<b>Immunogen</b>	Papain solubilized HLA-B7 antigen
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P01889</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">3106</a>    HLA-B    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	HLAB
<b>Specificity</b>	<p><b>Mouse anti Human HLA B7 antibody, clone BB7.1</b> recognizes the HLA B7 antigen and does not cross-react with HLA B27 or other related antigens. It can be used to distinguish true HLA B27 positives from false HLA B27 positives (i.e. HLA B7 positive) in the investigation of diseases such as ankylosing spondylitis and anterior uveitis. The major histocompatibility complex (MHC) is a cluster of genes that are important in the immune response to infections. In humans, this complex is referred to as the human leukocyte antigen (HLA) region. There are 3 major MHC class I proteins encoded by the HLA which are HLA A, HLA B and HLA C.</p> <p>The HLA B gene is part of the human HLA complex on chromosome 6 and there are a large number of variant alleles of this gene.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 1 x 10 <sup>6</sup> cells or 100ul whole blood
<b>References</b>	<ol style="list-style-type: none"> <li>1. Brodsky, F.M. <i>et al.</i> (1979) Monoclonal antibodies for analysis of the HLA system. <a href="#">Immunol Rev. 47: 3-61.</a></li> <li>2. Yoshino N <i>et al.</i> (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (<i>Macaca fascicularis</i>) by using anti-human cross-reactive antibodies. <a href="#">Exp Anim. 49 (2): 97-110.</a></li> <li>3. Anania, V.G. &amp; Coscoy, L. (2011) Palmitoylation of MIR2 is required for its function. <a href="#">J Virol. 85 (5): 2288-95.</a></li> <li>4. Bonaparte, M.I. and Barker, E. (2004) Killing of human immunodeficiency virus-infected primary T-cell blasts by autologous natural killer cells is dependent on the ability of the virus to alter the expression of major histocompatibility complex class I molecules. <a href="#">Blood. 104: 2087-94.</a></li> <li>5. Dellgren, C. <i>et al.</i> (2016) Low Constitutive Cell Surface Expression of HLA-B Is Caused by a Posttranslational Mechanism Involving Glu180 and Arg239. <a href="#">J Immunol. 197 (12): 4807-4816.</a></li> </ol>
<b>Storage</b>	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended.

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA986AMO">https://www.bio-rad-antibodies.com/SDS/MCA986AMO</a> 10041
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M360077:191028'

**Printed on 19 Oct 2023**