

Datasheet: MCA987

Description:	MOUSE ANTI HUMAN TCR ALPHA CHAIN
Specificity:	TCR ALPHA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	3A8
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry		•		
Immunohistology - Frozen	•			
Immunohistology - Paraffin			•	
ELISA			•	
Immunoprecipitation				
Western Blotting			•	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human	
Species Cross Reactivity	Does not react with:Mouse, Rat	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared from ascites by affinity chromatography	on Protein A
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃) 0.5% Bovine Serum Albumin	
Approx. Protein	IgG concentration 0.1 mg/ml	

Concentrations

External	Database
Links	

UniProt:

P01848 Related reagents

Entrez Gene:

28755 TRAC Related reagents

Synonyms	TCRA
RRID	AB_322432

Specificity

Mouse anti Human TCR alpha chain antibody, clone 3A8 reacts with the alpha chain constant region found on all alpha/beta TCR bearing T-lymphocytes. It reacts with 92-98% of all CD3 + peripheral blood lymphocytes.

Further Reading

- 1. Minami, Y. *et al.* (1987) Building a multichain receptor: synthesis, degradation, and assembly of the T-cell antigen receptor. <u>Proc Natl Acad Sci U S A. 84 (9): 2688-92.</u>
- 2. Koning, F. *et al.* (1988) The biosynthesis and assembly of T cell receptor alpha- and beta-chains with the CD3 complex. J Immunol. 140 (9): 3126-34.
- 3. van Dongen, J.J. *et al.* (1987) T cell receptor-CD3 complex during early T cell differentiation. Analysis of immature T cell acute lymphoblastic leukemias (T-ALL) at DNA, RNA, and cell membrane level. <u>J Immunol. 138 (4): 1260-9.</u>

Storage

Store at +4°C.

DO NOT FREEZE

This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA987 10041
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) RPE

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Goat Anti Mouse IgG (STAR76...)

Rabbit Anti Mouse IgG (STAR13...)

HRP

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Goat Anti Mouse IgG (STAR77...) HRP
Rabbit Anti Mouse IgG (STAR9...) FITC

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M389749:210806'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint