

Datasheet: MCA987T

Description:	MOUSE ANTI HUMAN TCR ALPHA CHAIN
Specificity:	TCR ALPHA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	3A8
Isotype:	IgG2a
Quantity:	50 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry		▪		
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Does not react with: Mouse, Rat
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared from ascites by affinity chromatography on Protein A
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃) 0.5% Bovine Serum Albumin
Approx. Protein	IgG concentration 0.1 mg/ml

Concentrations

External Database

Links

UniProt:

[P01848](#) [Related reagents](#)

Entrez Gene:

[28755](#) TRAC [Related reagents](#)

Synonyms

TCRA

RRID

AB_322682

Specificity

Mouse anti Human TCR alpha chain antibody, clone 3A8 reacts with the alpha chain constant region found on all alpha/beta TCR bearing T-lymphocytes. It reacts with 92-98% of all CD3 + peripheral blood lymphocytes.

Further Reading

1. Minami, Y. *et al.* (1987) Building a multichain receptor: synthesis, degradation, and assembly of the T-cell antigen receptor. [Proc Natl Acad Sci U S A. 84 \(9\): 2688-92.](#)
 2. Koning, F. *et al.* (1988) The biosynthesis and assembly of T cell receptor alpha- and beta-chains with the CD3 complex. [J Immunol. 140 \(9\): 3126-34.](#)
 3. van Dongen, J.J. *et al.* (1987) T cell receptor-CD3 complex during early T cell differentiation. Analysis of immature T cell acute lymphoblastic leukemias (T-ALL) at DNA, RNA, and cell membrane level. [J Immunol. 138 \(4\): 1260-9.](#)
-

Storage

Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10041 available at:
<https://www.bio-rad-antibodies.com/SDS/MCA987T>
10041

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

- Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
- Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
- Goat Anti Mouse IgG (STAR76...) [RPE](#)
- Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
- Goat Anti Mouse IgG (STAR70...) [FITC](#)
- Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@550](#),
[DyLight@650](#), [DyLight@680](#), [DyLight@800](#),
[FITC](#), [HRP](#)

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

Goat Anti Mouse IgG (STAR77...) [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M389750:210806'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)