

Datasheet: OBT0012 BATCH NUMBER 163426

Description:	MOUSE ANTI HUMAN TdT:FITC				
Specificity:	TdT				
Other names:	TERMINAL DEOXYNUCLEOTIDYL TRANSFERASE				
Format:	FITC				
Product Type:	Monoclonal Antibody Panel				
Clone:	Clone mixture				
Isotype:	Cocktail				
Quantity:	25 µg				

Product Details

Stabilisers

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal							
	communications from the originators. Please refer to references indicated for further							
		information. For general protocol recommendations, please visit <u>www.bio-</u>						
	· ·							
	rad-antibodies.com/protocols. Yes No Not Determined Suggested Dilution							
	Flow Cytometry (1)	Yes	NO	Not Determined	Suggested Dilution Neat			
	Immunohistology - Frozer	<u> </u>		•	INEAL			
	•••			•				
	Immunohistology - Paraffi	n		-				
	ELISA							
	Where this product has not been tested for use in a particular technique this does not							
	(1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code <u>BUF09</u>) is recommended for this purpose.							
Target Species	Human							
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid							
Max Ex/Em	Fluorophore	Excitation Max	(nm)	Emission Max (nm)				
	FITC	490		525				
Buffer Solution	Phosphate buffered sal	ine						

1% Bovine Serum Albumin

Approx. Protein Concentrations	IgG concentration 50.0 ug/ml
External Database Links	UniProt: P04053 Related reagents Entrez Gene: 1791 DNTT Related reagents
Synonyms	TDT
RRID	AB_609798
Specificity	Mouse anti Human TdT antibody panel is a mixture of 4 monoclonal antibodies specific for human terminal deoxynucleotidyl transferase (TdT). TdT is a DNA polymerase responsible for the catalysis of non-reversible addition of deoxynucleotides to the 3' end hydroxy groups of DNA. TdT levels are enhanced in all forms of acute lymphoblastic leukaemia (ALL) and in a significant number of chronic lymphoblastic leukaemia cases (CML). TdT is not expressed in the majority of myeloid leukaemias, non-Hodgkins lymphomas or mature lymphoid leukaemias.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
Flow Cytometry References	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul. 1. Wolgast, L.R. <i>et al.</i> (2011) Spectrin isoforms: differential expression in normal hematopoiesis and alterations in neoplastic bone marrow disorders. <u>Am J Clin Pathol.</u> <u>2011 Aug;136(2):300-8.</u>
	1. Wolgast, L.R. <i>et al.</i> (2011) Spectrin isoforms: differential expression in normal hematopoiesis and alterations in neoplastic bone marrow disorders. <u>Am J Clin Pathol.</u>
References	 Wolgast, L.R. <i>et al.</i> (2011) Spectrin isoforms: differential expression in normal hematopoiesis and alterations in neoplastic bone marrow disorders. <u>Am J Clin Pathol.</u> 2011 Aug;136(2):300-8. This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
References	 1. Wolgast, L.R. <i>et al.</i> (2011) Spectrin isoforms: differential expression in normal hematopoiesis and alterations in neoplastic bone marrow disorders. <u>Am J Clin Pathol.</u> 2011 Aug;136(2):300-8. This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be
References Storage	 Wolgast, L.R. <i>et al.</i> (2011) Spectrin isoforms: differential expression in normal hematopoiesis and alterations in neoplastic bone marrow disorders. <u>Am J Clin Pathol.</u> 2011 Aug;136(2):300-8. This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Related Products

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21				
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50	То			
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.comd a				
batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets									

'M404389:220822'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint