

Datasheet: OBT0012 BATCH NUMBER 168843

Description:	MOUSE ANTI HUMAN TdT:FITC
Specificity:	TdT
Other names:	TERMINAL DEOXYNUCLEOTIDYL TRANSFERASE
Format:	FITC
Product Type:	Monoclonal Antibody Panel
Clone:	Clone mixture
Isotype:	Cocktail
Quantity:	25 μg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	•			Neat
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	
ELISA			•	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code <u>BUF09</u>) is recommended for this purpose.

Target Species	Human			
Product Form	Purified IgG conju	ugated to Fluorescein Isotl	niocyanate Isomer 1	(FITC) - liquid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	FITC	490	525	
Preparation	Purified IgG from	ascites		
Buffer Solution	Phosphate buffer	ed saline		

Preservative Stabilisers	0.1% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 50.0 ug/ml
External Database Links	UniProt: P04053 Related reagents
	Entrez Gene: 1791 DNTT Related reagents
Synonyms	TDT
RRID	AB_609798
Specificity	Mouse anti Human TdT antibody panel is a mixture of 4 monoclonal antibodies specific for human terminal deoxynucleotidyl transferase (TdT). TdT is a DNA polymerase responsible for the catalysis of non-reversible addition of deoxynucleotides to the 3' end hydroxy groups of DNA. TdT levels are enhanced in all forms of acute lymphoblastic leukaemia (ALL) and in a significant number of chronic lymphoblastic leukaemia cases (CML). TdT is not expressed in the majority of myeloid leukaemias, non-Hodgkins lymphomas or mature lymphoid leukaemias.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	1. Wolgast, L.R. <i>et al.</i> (2011) Spectrin isoforms: differential expression in normal hematopoiesis and alterations in neoplastic bone marrow disorders. <u>Am J Clin Pathol.</u> 2011 Aug;136(2):300-8.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
Guarantee	Guaranteed until date of expiry. Please see product label.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/OBT0012 10041
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

<u>HUMAN SEROBLOCK (BUF070A)</u> <u>HUMAN SEROBLOCK (BUF070B)</u>

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M426421:231221'

Printed on 29 Aug 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint