# Datasheet: OBT1610 BATCH NUMBER 147466

Description:	MOUSE ANTI HUMAN NESTIN		
Specificity:	NESTIN		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	10C2		
lsotype:	lgG1		
Quantity:	0.1 mg		

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes	No	Not Determined	Suggested Dilution
	Immunohistology - Frozen	-			
	Immunohistology - Paraffin	-			1/50 - 1/200
	ELISA			•	
	Western Blotting	-			1/500 - 1/1000
	Where this product has n	ot been t	ested for u	use in a particular tech	nique this does not
	necessarily exclude its us	se in such	n procedui	es. Suggested working	g dilutions are given as
	a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.				or use in their own
Target Species	Human				
Species Cross Reactivity	Reacts with: Cynomolgus monkey Does not react with:Mouse, Rat <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.				
Product Form	Purified IgG - liquid				
Buffer Solution	Phosphate buffered saline				
Preservative	0.1% Sodium Azide (NaN <sub>3</sub> )				

#### Stabilisers

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	UniProt: <u>P48681</u> <u>Related reagents</u> Entrez Gene: <u>10763</u> NES <u>Related reagents</u>
RRID	AB_2151139
Specificity	<b>Mouse anti Human Nestin antibody, clone 10C2</b> recognizes nestin, a large intermediate filament protein originally identified by ( <u>Hockfield and McKay 1985</u> ). Nestin is predominately expressed in stem cells of the developing nervous system. Terminal differentiation is associated with a loss of nestin expression. Nestin expression has also been noted in other embryonic tissues, also in most Glioblastoma multiformes and many melanomas
Western Blotting	Suggested positive control: WI-38 (ATCC# CCL75) or U251 cell lysate.
References	<ol> <li>Hockfield, S. &amp; McKay, R.D. (1985) Identification of major cell classes in the developing mammalian nervous system. J Neurosci. 5 (12): 3310-28.</li> <li>Martínez-Navarrete, G.C. <i>et al.</i> (2008) Gradual morphogenesis of retinal neurons in the peripheral retinal margin of adult monkeys and humans J Comp Neurol. 511: 557-80.</li> <li>González-Garza, M.T. <i>et al.</i> (2013) Differentiation of CD133+ stem cells from amyotrophic lateral sclerosis patients into preneuron cells. Stem Cells Transl Med. 2 (2): 129-35.</li> <li>Aljammal, K. (2015) Combined Expression of Nestin and SPARC Identifies <i>In Situ</i> Tumor Cells in Astrocytic Tumors of all Grades J Cytol Histol. 6:313.</li> <li>Sterlacci, W. <i>et al.</i> (2014) Putative stem cell markers in non-small-cell lung cancer: a clinicopathologic characterization. J Thorac Oncol. 9 (1): 41-9.</li> <li>Nowak, A. <i>et al.</i> (2017) Nestin-positive microvessel density is an independent prognostic factor in breast cancer. Int J Oncol. Jun 26. [Epub ahead of print]</li> <li>Arranz, L. <i>et al.</i> (2016) Nestin(+) cells direct inflammatory cell migration in atherosclerosis. Nat Commun. 7: 12706.</li> <li>Forte, D. <i>et al.</i> (2020) Bone Marrow Mesenchymal Stem Cells Support Acute Myeloid Leukemia Bioenergetics and Enhance Antioxidant Defense and Escape from Chemotherapy. Cell Metab. 32 (5): 829-843.e9.</li> </ol>
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend

	microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/OBT1610 10040
Regulatory	For research purposes only

## **Related Products**

### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12)	<u>RPE</u>
Goat Anti Mouse IgG IgA IgM (STAR87	) <u>HRP</u>
Goat Anti Mouse IgG (STAR76)	<u>RPE</u>
Goat Anti Mouse IgG (STAR70)	FITC
Rabbit Anti Mouse IgG (STAR13)	HRP
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP
Rabbit Anti Mouse IgG (STAR9)	FITC
Goat Anti Mouse IgG (STAR77)	HRP
Goat Anti Mouse IgG (H/L) (STAR117)	<u>Alk. Phos.</u> , <u>DyLight®488</u> , <u>DyLight®550</u> ,
	DyLight®650, DyLight®680, DyLight®800,
	FITC, HRP

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M369335:200529'

#### Printed on 16 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint