Datasheet: PHP100 BATCH NUMBER 166350

Description:	RECOMBINANT HUMAN INTERLEUKIN-12
Name:	IL-12
Format:	Rec. Protein
Product Type:	Recombinant Protein
Quantity:	10 µg

Product Details

Applications	This product has been re derived from testing with communications from the information. For general rad-antibodies.com/proto	in our labo e originato protocol re	oratories, rs. Please	peer-reviewed publica e refer to references in	itions or personal ndicated for further
		Yes	No	Not Determined	Suggested Dilution
	ELISA	•			0.2 - 0.4 ng/well
	Western Blotting	•			1.5 - 3.0 ng/lane
	Functional Assays	-			0.1 - 10 ng/ml
	Where this protein has n	ot been te	sted for u	se in a particular tech	nique this does not
	necessarily exclude its u a guide only. It is recomr system using appropriate	mended th	at the use	er titrates the product f	
Target Species	Human				
Product Form	Purified recombinant pro	otein - Iyop	hilized		
Reconstitution	Reconstitute with 50ul st serum albumin is recomi Care should be taken du bottom of the vial. Bio-R Do not vortex.	mended. Iring recor	stitution a	as the protein may app	pear as a film at the
Preparation	Purified recombinant Hu cells.	man Interl	eukin-12 (expressed in Chinese	Hamster Ovary (CHO)
Source	CHO cells				
Buffer Solution	Phosphate buffered salir	ne			
Preservative	None present				

Stabilisers

Carrier Free	Yes
Endotoxin Level	< 0.1 ng/ug
Approx. Protein Concentrations	0.2 mg/ml after reconstitution
External Database Links	UniProt:
	P29460 Related reagents
	P29459 Related reagents
	Entrez Gene:
	3593 IL12B Related reagents
	<u>3592</u> IL12A <u>Related reagents</u>
Synonyms	NKSF1, NKSF2
Product Information	Recombinant human Interleukin-12 (IL-12) produced in CHO cells is a ~75 kDa glycoprotein comprised of disulfide-linked 35kDa (p35) and 40kDa (p40) subunits.
	IL-12 is produced by activated B lymphocytes and macrophages. The properties of IL-12 include the stimulation of growth of activated T lymphocytes, and actions in the development of Th1 CD4 +ve cells and CD8 +ve cytotoxic T cells. IL-12 also promotes activity and division of NK cells and promotes formation of TH1 cells producing &gamma-interferon.
Protein Molecular Weight	75kDa (503 Amino acid sequence)
Purity	>98% by SDS PAGE and HPLC analysis
ELISA	Recombinant human IL-12 may be used as a standard for ELISA applications with either anti-human IL-12:purified (AHP779) or anti-human IL-12:biotin conjugate (AHP779B).
Western Blotting	Recombinant human IL-12 may be used as the positive control for western blotting applications with either <u>anti-human IL-12:purified</u> (AHP779) or <u>anti-human IL-12:biotin</u> <u>conjugate</u> (AHP779B).
Storage	Prior to reconstitution store at -20°C. Following reconstitution store at -20°C.
	This product should be stored undiluted.
	Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the protein. Should this product contain a precipitate we recommend microcentrifugation before use.

I			•	heet documentation #10 htibodies.com/SDS/PHP	at:	
			s only			
	Tel: +1 800 265 7376 Worldwide Fax: +1 919 878 3751		Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739		Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50	
To find a b	Email: antibody_ atch/lot specif			Email: antibody_sales_uk@bio uct, please use our online s		Email: antibody_sales_de@bio-rad.co

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint