

Datasheet: PHP177

BATCH NUMBER 164857

Description:	RECOMBINANT HUMAN INSULIN-LIKE GROWTH FACTOR I
Name:	IGF-I
Other names:	INSULIN-LIKE GROWTH FACTOR I
Format:	Rec. Protein
Product Type:	Recombinant Protein
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Functional Assays	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified recombinant protein - lyophilized
Reconstitution	Reconstitute with 1.0 ml distilled water. Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.
Preparation	Purified recombinant IGF-I expressed in <i>E.coli</i>
Source	E.coli
Preservative Stabilisers	None present
Carrier Free	Yes
Endotoxin Level	< 1.0 EU/ug

Approx. Protein Concentrations	0.1 mg/ml
External Database Links	<p>UniProt: P05019 Related reagents</p> <p>Entrez Gene: 3479 IGF1 Related reagents</p>
Synonyms	IBP1
Product Information	<p>IGF-I (Insulin-like Growth Factor I), is a secreted mitogenic polypeptide and member of the insulin gene family, produced primarily by the liver, which effects a wide range of cells and plays a key role in cell proliferation and growth, inhibition of apoptosis, and is linked with tumour cell growth.</p> <p>IGF-I signals through binding to the type 1 insulin-like growth factor receptor (IGF-1R) and also to the insulin receptor (IR), resulting in the activation of the AKT signalling pathway. IGF-I production is stimulated by growth hormone (GH). and can be hindered by GH insensitivity, deficiency in GH receptors or a malfunction in the post GH receptor signalling pathway.</p>
Protein Molecular Weight	7.6 kDa
Activity	The ED ₅₀ was determined by a cell proliferation assay using FDC-P1 cells is < 2.0 ng/ml, corresponding to a specific activity of > 5 x 10 ⁵ units/mg.
Purity	>98% by SDS PAGE/HPLC
References	1. Zhang X <i>et al.</i> (2015) Wnt signaling regulates the stemness of lung cancer stem cells and its inhibitors exert anticancer effect on lung cancer SPC-A1 cells. Med Oncol. 32 (4): 462.
Storage	<p>Prior to reconstitution store at -20°C. After reconstitution store at -20°C. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Guarantee	Guaranteed for 3 months from the date of reconstitution or until the date of expiry, whichever comes first. Please see label for expiry date.
Health And Safety Information	Material Safety Datasheet documentation #10268 available at: https://www.bio-rad-antibodies.com/SDS/PHP177 10268
Regulatory	For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M415775:230113'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)