Datasheet: PHP177 BATCH NUMBER 164857

Description:	RECOMBINANT HUMAN INSULIN-LIKE GROWTH FACTOR I				
Name:	IGF-I				
Other names:	INSULIN-LIKE GROWTH FACTOR I				
Format:	Rec. Protein				
Product Type:	Recombinant Protein				
Quantity:	0.1 mg				

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>						
	rad-antibodies.com/protocols. Yes No Not Determined Suggested Dilution						
	Functional Assays						
	Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.						
Target Species	Human						
Product Form	Purified recombinant protein - lyophilized						
Reconstitution	Reconstitute with 1.0 ml distilled water. Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.						
Preparation	Purified recombinant IGF-I expressed in <i>E.coli</i>						
Source	E.coli						
Preservative Stabilisers	None present						
Carrier Free	Yes						
Endotoxin Level	< 1.0 EU/ug						

Approx. Protein Concentrations	0.1 mg/ml
External Database Links	UniProt: P05019 Related reagents Entrez Gene: 3479 IGF1 Related reagents
Synonyms	IBP1
Product Information	IGF-I (Insulin-like Growth Factor I), is a secreted mitogenic polypeptide and member of the insulin gene family, produced primarily by the liver, which effects a wide range of cells and plays a key role in cell proliferation and growth, inhibition of apoptosis, and is linked with tumour cell growth.
	IGF-I signals through binding to the type 1 insulin-like growth factor receptor (IGF-1R) and also to the insulin receptor (IR), resulting in the activation of the AKT signalling pathway. IGF-I production is stimulated by growth hormone (GH). and can be hindered by GH insensitivity, deficiency in GH receptors or a malfunction in the post GH receptor signalling pathway.
Protein Molecular Weight	7.6 kDa
Activity	The ED ₅₀ was determined by a cell proliferation assay using FDC-P1 cells is < 2.0 ng/ml, corresponding to a specific activity of > 5 x 10^5 units/mg.
Purity	>98% by SDS PAGE/HPLC
References	1. Zhang X <i>et al.</i> (2015) Wnt signaling regulates the stemness of lung cancer stem cells and its inhibitors exert anticancer effect on lung cancer SPC-A1 cells. <u>Med Oncol. 32 (4):</u> <u>462.</u>
Storage	Prior to reconstitution store at -20°C. After reconstitution store at -20°C. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	Guaranteed for 3 months from the date of reconstitution or until the date of expiry, whichever comes first. Please see label for expiry date.
Health And Safety Information	Material Safety Datasheet documentation #10268 available at: https://www.bio-rad-antibodies.com/SDS/PHP177 10268
Regulatory	For research purposes only

North & South	Tel: +1 800 265 7376	Vorldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M415775:230113'

Printed on 18 Nov 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint