Datasheet: PHP201 BATCH NUMBER 162289

Description:	RECOMBINANT HUMAN VISFATIN
Name:	VISFATIN
Other names:	NAMPT
Format:	Rec. Protein
Product Type:	Recombinant Protein
Quantity:	25 µg

Product Details

Applications	This product has been re derived from testing with communications from the information. For general rad-antibodies.com/proto	in our labo originato protocol re	oratories, p rs. Please	peer-reviewed publica refer to references in	tions or personal dicated for further
		Yes	No	Not Determined	Suggested Dilution
	Functional Assays Where this product has r	• ot been to	ested for u	se in a particular tech	prique this does not
	a guide only. It is recommonly system using appropriate	se in such nended th	procedure at the use	es. Suggested workin r titrates the product f	g dilutions are given as
Target Species	Human				
Product Form	Purified recombinant pro	tein - Iyop	hilized		
Reconstitution	Reconstitute with 0.25 ml distilled water. Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution. N.B. Due to solubility reasons, it is recommended that the solution be kept at low pH.				
Preparation	Purified recombinant visf	atin expre	ssed in <i>E.</i>	coli	
Buffer Solution	10mM HCI				
Preservative Stabilisers	None present				
Carrier Free	Yes				
Endotoxin Level	< 1.0 EU/ug				

Product Information	UniProt: P43490 Related reagents Entrez Gene: 10135 NAMPT Related reagents PBEF, PBEF1
Product Information	10135 NAMPT Related reagents
Product Information	PBEF, PBEF1
	Visfatin is a putative cytokine originally identified as PBEF (Pre-B cell-enhancing factor), subsequently determined to be a nicotinamide phosphoribosyltransferase (NAMPT) enzyme and member of the NAPRTase family, involved in catalyzing the biosynthesis of NAD from nicotinamide.
	Visfatin is highly expressed in bone marrow, muscle and liver tissue and increases in expression in response to endotoxin, in amniotic epithelial cells following lipopolysaccharide (LPS) stimulation, and during the activation of T and B cells, dendritic cells and macrophages, suggesting a role in inflammatory processes.
Protein Molecular Weight	52.5 kDa (465 amino acid residues)
	The ED $_{50}$ was determined by the dose-dependant proliferation of the RPMI 8226 cells. The expected ED $_{50}$ for this effect is 15.0-20.0 ng/ml.
Purity	>98% by SDS PAGE and HPLC analysis.
	Prior to reconstitution store at -20°C. After reconstitution store at -20°C. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Shoul this product contain a precipitate we recommend microcentrifugation before use.
	Guaranteed for 3 months from the date of reconstitution or until the date of expiry, whichever comes first. Please see label for expiry date.
Information	Material Safety Datasheet documentation #10268 available at: https://www.bio-rad-antibodies.com/SDS/PHP201 10268
Regulatory	For research purposes only

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint