

## Datasheet: PHP201 BATCH NUMBER 164851

Description:	RECOMBINANT HUMAN VISFATIN		
Name:	VISFATIN		
Other names:	NAMPT		
Format:	Rec. Protein		
Product Type:	Recombinant Protein		
Quantity:	25 μg		

## **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Functional Assays	•			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified recombinant protein - lyophilized
Reconstitution	Reconstitute with 0.25 ml distilled water. Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.  N.B. Due to solubility reasons, it is recommended that the solution be kept at low pH.
Preparation	Purified recombinant visfatin expressed in <i>E.coli</i>
Source	E.coli
Buffer Solution	10mM HCI
Preservative Stabilisers	None present
Carrier Free	Yes

Endotoxin Level	< 1.0 EU/ug
Approx. Protein Concentrations	Total protein concentration 0.1 mg/ml after reconstitution
External Database Links	UniProt: P43490 Related reagents  Entrez Gene:
	10135 NAMPT Related reagents
Synonyms	PBEF, PBEF1
Product Information	Visfatin is a putative cytokine originally identified as PBEF (Pre-B cell-enhancing factor), subsequently determined to be a nicotinamide phosphoribosyltransferase (NAMPT) enzyme and member of the NAPRTase family, involved in catalyzing the biosynthesis of NAD from nicotinamide.
	Visfatin is highly expressed in bone marrow, muscle and liver tissue and increases in expression in response to endotoxin, in amniotic epithelial cells following lipopolysaccharide (LPS) stimulation, and during the activation of T and B cells, dendritic cells and macrophages, suggesting a role in inflammatory processes.
Protein Molecular Weight	52.5 kDa (465 amino acid residues)
Activity	The ED $_{50}$ was determined by the dose-dependant proliferation of the RPMI 8226 cells. The expected ED $_{50}$ for this effect is 15.0-20.0 ng/ml.
Purity	>98% by SDS PAGE and HPLC analysis.
Storage	Prior to reconstitution store at -20°C.  After reconstitution store at -20°C.  Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	Guaranteed for 3 months from the date of reconstitution or until the date of expiry, whichever comes first. Please see label for expiry date.
Health And Safety Information	Material Safety Datasheet documentation #10268 available at: <a href="https://www.bio-rad-antibodies.com/SDS/PHP201">https://www.bio-rad-antibodies.com/SDS/PHP201</a> 10268

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

'M415642:230113'

Email: antibody\_sales\_de@bio-rad.com

## Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint