

Datasheet: PHP220

Description:	RECOMBINANT HUMAN M-CSF
Name:	M-CSF
Other names:	MACROPHAGE COLONY-STIMULATING FACTOR
Format:	Rec. Protein
Product Type:	Recombinant Protein
Quantity:	10 μg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
ELISA	•			0.2 - 0.4ng/well
Western Blotting	•			1.5 - 3.0ng/lane
Functional Assays	•			1.0 - 10.0ng/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human	
Product Form	Purified recombinant protein - lyophilized	
Reconstitution	Reconstitute with 0.1 ml distilled water. For extended storage recommended.	the addition of 0.1% BSA is
	Care should be taken during reconstitution as the protein may	• •
	bottom of the vial. Bio-Rad recommend that the vial is gently i	mixed after reconstitution.
Preparation	Purified recombinant M-CSF expressed in <i>E.coli</i>	
Source	E.coli	
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	None present	

Carrier Free	Yes
Endotoxin Level	< 1.0 EU/ug
Approx. Protein Concentrations	0.1 mg/ml after reconstitution
External Database Links	UniProt: P09603 Related reagents Entrez Gene: 1435 CSF1 Related reagents
Product Information	M-CSF (Macrophage colony-stimulating factor) is a potent lineage restricted haematopoietic growth factor and member of the CSF family of hormone-like glycoproteins, which plays a key role in the regulation of cellular differentiation, proliferation, and the survival of blood monocytes, tissue macrophages and their progenitor cells.
	Along with sRANKL/CD254/OPGL, M-CSF is essential for osteoclastogenesis, acting through the tyrosine kinase receptor c-fms (colony stimulating factor 1 receptor), and facilitates haematopoietic recovery following bone marrow transplantation.
Protein Molecular Weight	36.8 kDa (159 amino acid homodimer)
Activity	The ED ₅₀ , determined by the dose-dependent stimulation of the proliferation of M-NSF-60 cells, was \leq 1 ng/ml corresponding to a specific activity of \geq 1 x 10 ⁶ units/mg.
Purity	≥ 98% by SDS PAGE and HPLC
ELISA	PHP220 may be used in ELISA applications with <u>AHP1915</u> .
Western Blotting	PHP220 may be used under reducing or non-reducing western blotting conditions with <u>AHP1915</u> .
Storage	Prior to reconstitution store at -20°C. After reconstitution store at -20°C. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	Guaranteed for 3 months from the date of reconstitution or until the date of expiry, whichever comes first. Please see label for expiry date.
Health And Safety Information	Material Safety Datasheet documentation #10527 available at: https://www.bio-rad-antibodies.com/SDS/PHP220 10527

Regulatory For research purposes only

Related Products

Recommended Useful Reagents

RABBIT ANTI HUMAN M-CSF (AHP1915)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M420120:230706'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint