

Datasheet: PHP250

Description:	RECOMBINANT HUMAN CD54
Name:	CD54
Other names:	ICAM-1
Format:	Rec. Protein
Product Type:	Recombinant Protein
Quantity:	50 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
ELISA	▪			0.2 - 0.4ng/well
Western Blotting	▪			1.5 - 3.0ng/lane
Functional Assays	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified recombinant protein - lyophilized
Reconstitution	Reconstitute with 0.5 ml distilled water. Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.
Preparation	Purified recombinant CD54 expressed in CHO cells
Buffer Solution	10mM Sodium Phosphate
Preservative Stabilisers	None present
Carrier Free	Yes
Endotoxin Level	< 1.0 EU/ug
Approx. Protein Concentrations	0.1 mg/ml after reconstitution
External Database Links	UniProt:

Entrez Gene:

[3383](#) ICAM1 [Related reagents](#)

Product Information	CD54, otherwise known as ICAM-1, is a widely expressed type I integral membrane glycoprotein and member of the immunoglobulin superfamily, which acts as a ligand for the leukocyte integrin LFA-1 (CD11a/CD18) and Mac-1 integrins, and is the major receptor for human rhinovirus. CD54 is highly expressed on activated endothelial cells and plays a role in transendothelial cell migration to sites of acute inflammation, functioning primarily as an adhesion molecule between endothelial cells and leucocytes, following injury or stress. Levels of soluble CD54 in serum, can be potentially used as an indication of infection, inflammation and certain tumours, and as a target for immunosuppression in transplantation.
Protein Molecular Weight	49.5 kDa (453 amino acid residues)
Activity	CD54 supported the adhesion of more than 50% PMA-stimulated HSB2 cells to plates coated at a CD54 concentration of 25ng/ml.
Purity	>98% by SDS PAGE and HPLC analysis
ELISA	PHP250 may be used as a standard in ELISA applications involving AHP2183 or AHP2183B .
Western Blotting	PHP250 may be used as a positive control for western blotting applications involving AHP2183 or AHP2183B .
Storage	Prior to reconstitution store at +4°C. After reconstitution store at -20°C. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	3 months from date of reconstitution
Health And Safety Information	Material Safety Datasheet documentation #10345 available at: 10345: https://www.bio-rad-antibodies.com/uploads/MSDS/10345.pdf
Regulatory	For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M362384:200501'

Printed on 11 Aug 2020