

Datasheet: PHP251

Description:	RECOMBINANT HUMAN CD141
Name:	CD141
Other names:	THROMBOMODULIN
Format:	Rec. Protein
Product Type:	Recombinant Protein
Quantity:	10 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Functional Assays	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified recombinant protein - lyophilized
Reconstitution	Reconstitute with 0.1ml distilled water Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution. Further may be prepared in a buffer containing 0,1% BSA.
Preparation	Purified recombinant CD141 expressed in HEK293 cells
Source	HEK-293
Buffer Solution	10mM Sodium Phosphate
Preservative Stabilisers	None present
Carrier Free	Yes

Endotoxin Level	< 1.0 EU/ug
Approx. Protein Concentrations	0.1mg/ml after reconstitution
External Database Links	<p>UniProt: P07204 Related reagents</p> <p>Entrez Gene: 7056 THBD Related reagents</p>
Synonyms	THRM
Product Information	<p>Recombinant Human CD141 antigen produced in HEK293 cells represents full length human CD141, otherwise known as Thrombomodulin, is a transmembrane glycoprotein specifically synthesized by endothelial cells, which plays a critical role in the activation/conversion of protein C into protein Ca, through the formation of a complex with the coagulation factor Thrombin. The subsequent cleavage and deactivation of the coagulation factors Va and factor VIIIa by protein Ca, acts to reduce Thrombin production and hence prevent further coagulation.</p> <p>Plasma levels of CD141 are increased in diabetes mellitus and SLE, and Thrombomodulin can correlate with the pathogenesis of the cardiovascular diseases thrombosis and atherosclerosis.</p> <p>Recombinant soluble Thrombomodulin contains 491 amino acid residues, including the extracellular domain of Thrombomodulin.</p>
Activity	Measured by its ability to activate protein C induced cleavage of the chromogenic substrate, BOC-Asp-Pro Arg-AMC in the presence of Thrombin. The specific activity is >500 pmoles/min/ug.
Purity	>98% by SDS PAGE and HPLC analysis
Storage	<p>Prior to reconstitution store at +4°C.</p> <p>After reconstitution store at -20°C.</p> <p>Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Guarantee	3 months from date of reconstitution
Health And Safety Information	<p>Material Safety Datasheet documentation #10527 available at: https://www.bio-rad-antibodies.com/SDS/PHP25110527</p>
Regulatory	For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com	To find a
----------------------------------	---	------------------	---	---------------	---	-----------

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M420128:230706'

Printed on 13 Aug 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)