

Datasheet: PHP295

BATCH NUMBER 100005637

| Description: | RECOMBINANT HUMAN CCL17 |
|---------------|-------------------------|
| Name: | CCL17 |
| Format: | Rec. Protein |
| Product Type: | Recombinant Protein |
| Quantity: | 10 µg |

UniProt:

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|-------|-----|----|----------------|--------------------|
| ELISA | | | | |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species | Human |
|-----------------------------------|--|
| Product Form | Purified recombinant protein - lyophilized |
| Reconstitution | Centrifuge vial prior to reconstitution. Reconstitute to 100 μ g/ml by adding 100 μ l ddH ₂ O. Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution. Do not vortex. |
| Preparation | Recombinant protein expressed in <i>E. coli</i> and purified by ion exchange chromatography |
| Buffer Solution | 50 mM Tris-HCl, pH 8.0, 0.15 M Sodium Chloride |
| Preservative Stabilisers | 1.0% Trehalose |
| Approx. Protein Concentrations | 100 μg/ml after reconstitution |
| External Database | |

| Links | Q92583 Related reagents |
|--------------------------------------|---|
| | Entrez Gene: 6361 CCL17 Related reagents |
| Synonyms | SCYA17, TARC |
| Product Information | Recombinant Human CCL17 |
| | CCL17 also known as TARC and SCYA17 is a C-C chemokine highly expressed by the thymus and other cell types, including fibroblasts (<u>Li et al. 2019</u>). CCL17 induces and enhances a broad spectrum of immune reactions, including some contact hypersensitivity reactions, allograft rejections, inflammatory bowel disease, atopic dermatitis and atherosclerosis (<u>Scheu et al. 2017</u>). Therapeutic use of antibodies to neutralize CCL17 can block influx of T cells and ameliorate the impact of various diseases (<u>Teplyakov et al. 2017</u>). Upregulation of CCL17 has been reported in breast cancer, and predicts a decrease in overall survival for patients with breast cancer (<u>Li et al. 2019</u>). |
| | This recombinant Human CCL17 can be used to create a standard curve in a sandwich ELISA with purified Mouse anti Human CCL17 antibody, clone I02-2A4 (MCA6180GA) as the capture antibody and biotinylated Mouse anti Human CCL17 antibody, clone I11-2A1 (MCA6181B) as the detection antibody. |
| Protein Molecular | 0.445 |
| Weight | 8.1 kDa |
| Weight Purity | 8.1 kDa ≥90 % determined by coomassie blue staining of a reducing SDS-PAGE gel |
| Purity | |
| Purity | ≥90 % determined by coomassie blue staining of a reducing SDS-PAGE gel MARGTNVGRE CCLEYFKGAI PLRKLKTWYQ TSEDCSRDAI VFVTVQGRAI |
| Purity Amino Acid Sequence | ≥90 % determined by coomassie blue staining of a reducing SDS-PAGE gel MARGTNVGRE CCLEYFKGAI PLRKLKTWYQ TSEDCSRDAI VFVTVQGRAI CSDPNNKRVK NAVKYLQSLE RS |
| Purity Amino Acid Sequence | ≥90 % determined by coomassie blue staining of a reducing SDS-PAGE gel MARGTNVGRE CCLEYFKGAI PLRKLKTWYQ TSEDCSRDAI VFVTVQGRAI CSDPNNKRVK NAVKYLQSLE RS Prior to reconstitution store at -70°C. Following reconstitution store at -80°C. |
| Purity Amino Acid Sequence | ≥90 % determined by coomassie blue staining of a reducing SDS-PAGE gel MARGTNVGRE CCLEYFKGAI PLRKLKTWYQ TSEDCSRDAI VFVTVQGRAI CSDPNNKRVK NAVKYLQSLE RS Prior to reconstitution store at -70°C. Following reconstitution store at -80°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the protein. Should this product contain a precipitate we recommended. |
| Purity Amino Acid Sequence Storage | ≥90 % determined by coomassie blue staining of a reducing SDS-PAGE gel MARGTNVGRE CCLEYFKGAI PLRKLKTWYQ TSEDCSRDAI VFVTVQGRAI CSDPNNKRVK NAVKYLQSLE RS Prior to reconstitution store at -70°C. Following reconstitution store at -80°C. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the protein. Should this product contain a precipitate we recommend microcentrifugation before use. Guaranteed for 3 months from the date of reconstitution or until the date of expiry, |

Related Products

Recommended Useful Reagents

MOUSE ANTI HUMAN CCL17 (MCA6180GA)
MOUSE ANTI HUMAN CCL17:Biotin (MCA6181B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M380420:210511'

Printed on 18 Nov 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint