

Datasheet: PMP48

BATCH NUMBER 180220

Description:	MOUSE IgG1 KAPPA
Name:	IgG1 KAPPA
Format:	Purified
Product Type:	Purified Protein
Clone:	MADNP-1
Isotype:	IgG1
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
ELISA	▪			0.6 ug/ml - 1.25 ug/ml
Gel Super Shift Assays	▪			

Where this protein has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the protein for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified monoclonal IgG1 - liquid
Preparation	Purified IgG1 prepared by affinity chromatography from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	UniProt: P01837 Related reagents

[P01869](#) [Related reagents](#)

[P01868](#) [Related reagents](#)

Entrez Gene:

[16071](#) Igk-C [Related reagents](#)

[16017](#) Ighg1 [Related reagents](#)

[16017](#) Ighg1 [Related reagents](#)

Synonyms Igh-4

RRID AB_324554

Product Information **Mouse IgG1 Kappa**

Purity Single band by agarose electrophoresis and immunoelectrophoresis.

References 1. Kiesler, P. *et al.* (2009) An allergy-associated polymorphism in a novel regulatory element enhances IL13 expression. [Hum Mol Genet. 18: 4513-20.](#)
2. Brown, J.K. *et al.* (2011) Human Fallopian tube epithelium constitutively expresses integrin endometrial receptivity markers: no evidence for a tubal implantation window. [Mol Hum Reprod. 18: 111-20.](#)
3. Laffont, S. *et al.* (2017) Androgen signaling negatively controls group 2 innate lymphoid cells. [J Exp Med. May 8. pii: jem.20161807. \[Epub ahead of print\]](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the protein. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at:
<https://www.bio-rad-antibodies.com/SDS/PMP48>
10040

Regulatory For research purposes only

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M369569:200529'

Printed on 19 Jan 2024
