

Datasheet: PRABP01

Description:	PURIFIED RABBIT IgG
Name:	IgG
Format:	Purified
Product Type:	Purified Protein
Isotype:	Polyclonal IgG
Quantity:	10 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
ELISA	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Rabbit
Product Form	Purified polyclonal rabbit IgG - liquid
Preparation	Purified IgG prepared by ion exchange chromatography from normal rabbit serum.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 2 mg/ml
External Database Links	<p>UniProt: P01870 Related reagents</p> <p>Entrez Gene: 100009097 LOC100009097 Related reagents</p>

RRID AB_321631

Product Information Rabbit immunoglobulin G is intended as a negative control reagent for use alongside specific rabbit IgG primary antibodies. It has been tested in flow cytometry on human tissues and demonstrates negligible binding. Purified Rabbit immunoglobulin G may also be used as a standard for rabbit IgG in ELISA ([Blakney et al. 2021](#)).

References

1. Khalifeh, M.S. *et al.* (2010) Investigation of the role of tumour necrosis factor- α , interleukin-1 β , interleukin-10, nitric oxide and rheumatoid factor-immunoglobulin M in a rat model of arthritis. [Lab Anim. 44: 143-9.](#)
2. Chang, J. *et al.* (2014) Hypoxia modulates the phenotype of osteoblasts isolated from knee osteoarthritis patients, leading to undermineralized bone nodule formation. [Arthritis Rheumatol. 66 \(7\): 1789-99.](#)
3. Blakney, A.K. *et al.* (2021) Innate Inhibiting Proteins Enhance Expression and Immunogenicity of Self-Amplifying RNA. [Mol Ther. 29 \(3\): 1174-85.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10258 available at: 10258: <https://www.bio-rad-antibodies.com/uploads/MSDS/10258.pdf>

Regulatory For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M389770:210806'

Printed on 21 Mar 2022

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)