

## Datasheet: PRABP01

<b>Description:</b>	PURIFIED RABBIT IgG
<b>Name:</b>	IgG
<b>Format:</b>	Purified
<b>Product Type:</b>	Purified Protein
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	10 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
ELISA	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Rabbit
<b>Product Form</b>	Purified polyclonal rabbit IgG - liquid
<b>Preparation</b>	Purified IgG prepared by ion exchange chromatography from normal rabbit serum.
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	<0.1% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 2 mg/ml
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P01870</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">100009097</a>    LOC100009097    <a href="#">Related reagents</a></p>

RRID AB\_321631

---

**Product Information** **Rabbit immunoglobulin G** is intended as a negative control reagent for use alongside specific rabbit IgG primary antibodies. It has been tested in flow cytometry on human tissues and demonstrates negligible binding. Purified Rabbit immunoglobulin G may also be used as a standard for rabbit IgG in ELISA ([Blakney \*et al.\* 2021](#)).

---

**References**

1. Khalifeh, M.S. *et al.* (2010) Investigation of the role of tumour necrosis factor- $\alpha$ , interleukin-1 $\beta$ , interleukin-10, nitric oxide and rheumatoid factor-immunoglobulin M in a rat model of arthritis. [Lab Anim. 44: 143-9.](#)
2. Chang, J. *et al.* (2014) Hypoxia modulates the phenotype of osteoblasts isolated from knee osteoarthritis patients, leading to undermineralized bone nodule formation. [Arthritis Rheumatol. 66 \(7\): 1789-99.](#)
3. Blakney, A.K. *et al.* (2021) Innate Inhibiting Proteins Enhance Expression and Immunogenicity of Self-Amplifying RNA. [Mol Ther. 29 \(3\): 1174-85.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10258 available at: <https://www.bio-rad-antibodies.com/SDS/PRABP0110258>

---

**Regulatory** For research purposes only

---

**North & South America** Tel: +1 800 265 7376  
Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide** Tel: +44 (0)1865 852 700  
Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe** Tel: +49 (0) 89 8090 95 21  
Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M389770:210806'

Printed on 19 Jan 2024

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)