

Datasheet: PRABP01F BATCH NUMBER 149809

Description:	PURIFIED RABBIT IgG:FITC		
Name:	IgG		
Format:	FITC		
Product Type:	Purified Protein		
Isotype:	Polyclonal IgG		
Quantity:	0.1 mg		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				*
Immunohistology - Frozen				
Immunohistology - Paraffin				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. * It is recommended that the user dilutes the antibody for use in their own system to a concentration equivalent to their test reagent.

Target Species	Rabbit				
Product Form	Purified polyclonal ra	abbit IgG conjugated to	Fluorescein Isothio		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm		
	FITC	490	525		
Preparation	Purified IgG prepared by ion exchange chromatography from normal rabbit serum				
Buffer Solution	Phosphate buffered saline				
Preservative	0.09% Sodium Azide	•			
Stabilisers	1% Bovine Serum Albumin				
Approx. Protein Concentrations	IgG concentration 0.	1 mg/ml			

External Database Links	UniProt: P01870 Related reagents Entrez Gene: 100009097 LOC100009097 Related reagents				
RRID	AB_324021				
Product Information	This product is designed as a negative control reagent for use alongside specific rabbit IgG primary antibodies. It has been tested in flow cytometry on human tissues and demonstrates negligible binding.				
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul of whole blood				
References	 Khalifeh, M.S. <i>et al.</i> (2010) Investigation of the role of tumour necrosis factor-{alpha}, interleukin-1{beta}, interleukin-10, nitric oxide and rheumatoid factor-immunoglobulin M in a rat model of arthritis. <u>Lab Anim. 44: 143-9.</u> Chang, J. <i>et al.</i> (2014) Hypoxia modulates the phenotype of osteoblasts isolated from knee osteoarthritis patients, leading to undermineralized bone nodule formation. <u>Arthritis Rheumatol. 66 (7): 1789-99.</u> 				
Storage	Store at +4°C or at -20°C if preferred.				
	This product should be stored undiluted. This product is photosensitive and should be protected from light.				
	Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.				
Guarantee	12 months from date of despatch				
Health And Safety Information	Material Safety Datasheet documentation #10259 available at: https://www.bio-rad-antibodies.com/SDS/PRABP01F 10259				
Regulatory	For research purposes only				

North & South Tel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700 Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 America Email: antibody_sales_us@bio-rad.com

Europe

Fax: +49 (0) 89 8090 95 50

Tel: +49 (0) 89 8090 95 21

Email: antibody_sales_uk@bio-rad.com Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M369579:200529'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint