

Datasheet: PRP07A BATCH NUMBER 158754

Description:	cription: RAT IgE KAPPA	
Name:	IgE KAPPA	
Format:	Purified	
Product Type:	Purified Protein	
Clone:	IR162	
Isotype:	IgE	
Quantity:	0.1 mg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
ELISA				0.1ug/ml - 1.25ug/ml

Where this protein has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the protein for use in their own system using appropriate positive/negative controls.

Target Species	Rat	
Product Form	IgE purified (myeloma IR162) - liquid	
Preparation	Purified IgE prepared by affinity chromatography from tissue of	culture supernatant
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.1% Sodium Azide	
Approx. Protein Concentrations	IgE concentration 1.0 mg/ml	
External Database Links	UniProt:	

P01855 Related reagents

Entrez Gene:

299351 Ighe Related reagents

DDID	AD 400077
RRID	AB_482877
Specificity	Rat IgE is generated by a spontaneous immunocytoma isolated from a LOU/Wsl rat (<u>Bazin 1974</u> , <u>Bazin & Beckers 1974</u>). The determinant recognized by IgE clone IR162 appears to be shared by several helminthic parasites, including <i>Anisakis simplex</i> , <i>Clonorchis sinensis</i> and <i>Toxocara canis</i> (<u>Cho et al. 2000</u>).
Purity	Single band by agarose eletrophoresis (5) and immunoelectrophoresis (6)
References	 Bazin, H. <i>et al.</i> (1973) Brief communication: Transplantable immunoglobulin-secreting tumors in rats. V. Monoclonal immunoglobulins secreted by 250 ileocecal immunocytomas in LOU-Wsl rats. J. Natl. Cancer Inst. 51: 1359-1361. Bazin, H. <i>et al.</i> (1974) Transplantable immunoglobulin-secreting tumours in rats. IV. Sixty-three IgE-secreting immunocytoma tumours. Immunology. 26: 713-723. Bazin, H. and Beckers, A. (1974) IgE- Myelomas in Rats Nobel Symposium 33: Molecular and Biological Aspects of the Acute Allergic Reaction. Zeuthen, J. <i>et al.</i> (1980) IgE production in vitro by two rat myeloma tumor lines (IR2 and IR162). Cell Differ. 9 (3): 137-51. Bazin, H. <i>et al.</i> (1984) Rat monoclonal antibodies. II. A rapid and efficient method of purification from ascitic fluid or serum. J. Immunol. Methods. 71: 9-16. Bazin, H. J. <i>et al.</i> (1984) Rat monoclonal antibodies. I. Rapid purification from in vitro culture supernatants. Immunol. Methods. 66: 261-269. Bazin, H. (Ed) (1990) Rat Hybridomas and Rat Monoclonal Antibodies CRC Press, Boca Raton, Florida, USA. Tulinská J <i>et al.</i> (2018) Humoral and cellular immune response in Wistar Han RCC rats fed two genetically modified maize MON810 varieties for 90 days (EU 7th Framework Programme project GRACE). Arch Toxicol. 92 (7): 2385-99. Masuda, A. <i>et al.</i> (2018) Body Protein Reserves Sustain Maternal Performance in Early Lactation but Dietary Protein Is Necessary to Maintain Performance and Immune Responses to <i>Nippostrongylus brasiliensis</i>. in Lactating Rats. J Nutr. 148 (10): 1638-46.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in
	frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10335 available at: https://www.bio-rad-antibodies.com/SDS/PRP07A 10335

Regulatory For research purposes only

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385951:210513'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint