

Datasheet: STAR104

BATCH NUMBER 172171

Description:	GOAT F(ab') ₂ ANTI HAMSTER IgG
Specificity:	IgG
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.4 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/1000 - 1/2000

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Hamster

Product Form F(ab')₂ fragment of purified IgG - liquid

Antiserum Preparation Antisera to hamster IgG were raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography. F(ab')₂ fragments were prepared by pepsin digestion.

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.09% Sodium Azide

Approx. Protein Concentrations IgG concentration 0.8 mg/ml

Immunogen	Hamster IgG.
RRID	AB_323008
Specificity	Goat F(ab')₂ anti Hamster IgG antibody recognizes Golden Syrian and Armenian hamster IgG. Goat F(ab') ₂ anti Hamster IgG antibody has been adsorbed against both mouse and rat immunoglobulins to minimise cross-reactivity.
References	<p>1. Osorio, Y. <i>et al.</i> (2011) Identification of small molecule lead compounds for visceral leishmaniasis using a novel <i>ex vivo</i> splenic explant model system. PLoS Negl Trop Dis. 5 (2): e962.</p> <p>2. Bouma, G. <i>et al.</i> (2011) Cytoskeletal remodeling mediated by WASp in dendritic cells is necessary for normal immune synapse formation and T-cell priming. Blood. 118 (9): 2492-501.</p>
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/STAR104
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M428610:240301'

Printed on 29 Jan 2026