

Datasheet: STAR104D650 BATCH NUMBER 152512

Description:	GOAT F(ab')2 ANTI HAMSTER IgG:DyLight®650	
Specificity:	IgG	
Format:	DyLight®650	
Product Type:	Polyclonal Antibody	
Isotype:	Polyclonal IgG	
Quantity:	0.1 mg	

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>				
	rad-antibodies.com/protocols.				
	Flow Outomotry	Yes	No	Not Determined	Suggested Dilution 1/50 - 1/200
	Flow Cytometry				
					1/50 - 1/200
	Where this antibody ha		•	•	
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.				
Target Species	Hamster				
Product Form	F(ab') ₂ fragment of purified IgG conjugated to DyLight [®] 650 - liquid				
Max Ex/Em	Fluorophore	Excitation M	/lax (nm)	Emission Max (nm)	
	Dylight®650	654		673	
Antiserum Preparation	Antisera to hamster IgG were raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography.				
Buffer Solution	Phosphate buffered saline				
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)				
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml				

Immunogen	Hamster IgG.					
Specificity	Goat F(ab')2 anti Hamster IgG antibody recognizes Golden Syrian and Armenian hamster IgG. Goat F(ab')2 anti Hamster IgG antibody has been adsorbed against both mouse and rat immunoglobulins to minimise cross-reactivity.					
Flow Cytometry	Use 50ul of the suggested working dilution to label 1×10^6 cells in 100ul					
References	 Osorio, Y. <i>et al.</i> (2011) Identification of small molecule lead compounds for visceral leishmaniasis using a novel <i>ex vivo</i> splenic explant model system. <u>PLoS Negl Trop Dis. 5</u> (2): e962. Bouma, G. <i>et al.</i> (2011) Cytoskeletal remodeling mediated by WASp in dendritic cells is necessary for normal immune synapse formation and T-cell priming. <u>Blood. 118 (9):</u> 2492-501. 					
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.					
Guarantee	12 months from date of despatch					
Acknowledgements	$DyLight^{ extsf{B}}$ is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries					
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/STAR104D650 10040					
Regulatory	For research purposes only					
North & South Tel: +1 800 265 7 America Fax: +1 919 878 Email: antibody_s						
To find a batch/lot specif	ic datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M409075:221017'					

Printed on 29 Aug 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint