

Datasheet: STAR104F

Description:	GOAT F(ab')2 ANTI HAMSTER IgG:FITC
Specificity:	IgG
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.4 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				1/100 - 1/200
Immunohistology - Frozen				
Immunohistology - Paraffin				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Hamster			
F(ab')2 fragment o	of purified IgG conjugated	to Fluorescein Isothic	ocyanate Isomer 1 (FITC)
Fluorophore FITC	Excitation Max (nm) 490	Emission Max (nm)	
	F(ab')2 fragment of a liquid	F(ab')2 fragment of purified IgG conjugated - liquid Fluorophore Excitation Max (nm)	F(ab')2 fragment of purified IgG conjugated to Fluorescein Isothic - liquid Fluorophore Excitation Max (nm) Emission Max (nm)

Antiserum Preparation Antisera to hamster IgG were raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 0.5% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.8 mg/ml

Immunogen	Hamster IgG.
RRID	AB_323007
Specificity	Goat F(ab')2 anti Hamster IgG antibody recognizes Golden Syrian and Armenian hamster IgG. Goat F(ab')2 anti Hamster IgG antibody has been adsorbed against both mouse and rat immunoglobulins to minimise cross-reactivity.
Flow Cytometry	Use 50ul of the suggested working dilution to label 1 x 10^6 cells in 100ul.
References	1. Osorio, Y. <i>et al.</i> (2011) Identification of small molecule lead compounds for visceral leishmaniasis using a novel <i>ex vivo</i> splenic explant model system. <u>PLoS Negl Trop Dis. 5</u> (2): e962.
	2. Bouma, G. <i>et al.</i> (2011) Cytoskeletal remodeling mediated by WASp in dendritic cells is necessary for normal immune synapse formation and T-cell priming. <u>Blood. 118 (9): 2492-501.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/STAR104F 10041
Regulatory	For research purposes only

North & South America

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M428616:240301'

Printed on 29 Aug 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint