

Datasheet: STAR106

Description:	GOAT ANTI HUMAN IgG GAMMA CHAIN
Specificity:	IgG GAMMA CHAIN
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.8 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			1ug/ml - 5ug/ml
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Human

Product Form Purified IgG - liquid

Antiserum Preparation Antisera to human IgG were raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.09% Sodium Azide

Approx. Protein Concentrations IgG concentration 0.8 mg/ml

Immunogen Human IgG.

External Database **UniProt:**

Links

[P01857](#) [Related reagents](#)
[P01859](#) [Related reagents](#)
[P01860](#) [Related reagents](#)
[P01861](#) [Related reagents](#)

Entrez Gene:

[3500](#) IGHG1 [Related reagents](#)
[3501](#) IGHG2 [Related reagents](#)
[3502](#) IGHG3 [Related reagents](#)
[3503](#) IGHG4 [Related reagents](#)

RRID

AB_322999

Specificity

Goat anti Human IgG gamma chain antibody recognizes human IgG and demonstrates <1% cross-reactivity with other human immunoglobulin classes or subclasses. Goat anti Human IgG gamma Chain antibody recognizes the Fc region of IgG, and has been adsorbed against mouse immunoglobulins to reduce species cross-reactivity.

References

1. Nielsen, L.S. *et al.* (2010) Single-batch production of recombinant human polyclonal antibodies. [Mol Biotechnol. 45 \(3\): 257-66.](#)
2. Nordström, T. *et al.* (2011) Human Siglec-5 inhibitory receptor and immunoglobulin A (IgA) have separate binding sites in streptococcal beta protein. [J Biol Chem. 286 \(39\): 33981-91.](#)
3. Hanitsch, L.G. *et al.* (2015) Late-Onset Disseminated Mycobacterium avium intracellulare Complex Infection (MAC), Cerebral Toxoplasmosis and *Salmonella* Sepsis in a German Caucasian Patient with Unusual Anti-Interferon-Gamma IgG1 Autoantibodies. [J Clin Immunol. 35 \(4\): 361-5.](#)
4. Zheng, L. *et al.* (2015) Streptococcal 5'-nucleotidase A (S5nA), a novel group A Streptococcal virulence factor that facilitates immune evasion. [J Biol Chem. pii: jbc.M115.677443.](#)
5. Le, Y. *et al.* (2016) Immunogenicity of Murine mPEG-Red Blood Cells and the Risk of Anti-PEG Antibodies in Human Blood Donors. [Exp Hematol. Nov 15. pii: S0301-472X\(16\)30702-0. \[Epub ahead of print\]](#)
6. Zheng, L. *et al.* (2015) Streptococcal 5'-Nucleotidase A (S5nA), a Novel *Streptococcus pyogenes* Virulence Factor That Facilitates Immune Evasion. [J Biol Chem. 290 \(52\): 31126-37.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety

Material Safety Datasheet documentation #10040 available at:

Information 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M381994:210512'

Printed on 21 Mar 2022

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)