

## Datasheet: STAR117A

**BATCH NUMBER 162670**

<b>Description:</b>	GOAT ANTI MOUSE IgG (H/L):Alk. Phos. (MULTI SPECIES ADSORBED)
<b>Specificity:</b>	IgG (H/L)
<b>Format:</b>	Alk. Phos.
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	0.5 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/5000 - 1/50000
Immunoprecipitation			▪	
Western Blotting	▪			1/2500 - 1/25000

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

#### Target Species

Mouse

#### Product Form

Purified IgG conjugated to Alkaline Phosphatase - liquid

#### Antiserum Preparation

Antisera to mouse IgG were raised by repeated immunisations of goats with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

#### Buffer Solution

50 mM HEPES pH 7.1  
0.1M NaCl  
1mM MgCl<sub>2</sub>  
0.1mM ZnCl<sub>2</sub>

#### Preservative Stabilisers

0.09% Sodium Azide

0.2% Bovine Serum Albumin

---

**Approx. Protein Concentrations** IgG concentration 0.5 mg/ml

---

**Immunogen** Whole mouse IgG

---

**External Database Links**

**UniProt:**

[P01837](#) [Related reagents](#)  
[P01869](#) [Related reagents](#)  
[P01867](#) [Related reagents](#)  
[P01864](#) [Related reagents](#)  
[P01843](#) [Related reagents](#)  
[P01865](#) [Related reagents](#)  
[P01844](#) [Related reagents](#)  
[P01868](#) [Related reagents](#)  
[P01724](#) [Related reagents](#)  
[P03987](#) [Related reagents](#)  
[P01863](#) [Related reagents](#)  
[P01845](#) [Related reagents](#)

**Entrez Gene:**

[16071](#) Igk-C [Related reagents](#)  
[16017](#) Ighg1 [Related reagents](#)  
[16016](#) Ighg2b [Related reagents](#)  
[380793](#) Igh-1a [Related reagents](#)  
[380793](#) Igh-1a [Related reagents](#)  
[433053](#) LOC433053 [Related reagents](#)  
[16017](#) Ighg1 [Related reagents](#)  
[16142](#) Iglv1 [Related reagents](#)  
[110786](#) Iglc2 [Related reagents](#)  
[110787](#) Iglc3 [Related reagents](#)  
[380793](#) Igh-1a [Related reagents](#)  
[380795](#) AI324046 [Related reagents](#)

---

**Synonyms** Igh-4

---

**RRID** AB\_324391

---

**Specificity** **Goat anti Mouse IgG antibody** recognizes mouse IgG and light chains common to other mouse immunoglobulin classes.

Goat anti Mouse IgG has been cross-adsorbed using human, bovine, porcine, equine, lapine and chicken immunoabsorbants to remove cross-reactive antibodies. Less than 0.1% cross reactivity was detected to human, bovine, porcine, equine, caprine, lapine and

chicken IgG by immunoelectrophoresis and ELISA.

Goat anti Mouse IgG antibody is highly recommended for use as a secondary antibody with human and veterinary samples. Goat anti Mouse IgG antibody has been used successfully as a secondary detection reagent in combination with mouse clone [CC327](#) for the detection of TNF $\alpha$  and mouse clone [8M6](#) for the detection of interleukin-8 in bovine respiratory syncytial virus infected, neonatal ovine lung tissue by immunohistochemistry ([Redondo et al. 2013](#)).

---

## References

1. Abdala-Valencia, H. *et al.* (2012) Vitamin E isoforms differentially regulate intercellular adhesion molecule-1 activation of PKC $\alpha$  in human microvascular endothelial cells. [PLoS One. 7: e41054.](#)
2. Redondo, E. *et al.* (2014) Induction of interleukin-8 and interleukin-12 in neonatal ovine lung following experimental inoculation of bovine respiratory syncytial virus. [J Comp Pathol. 150 \(4\): 434-48.](#)
3. Banerjee, K. *et al.* (2012) Occluding the mannose moieties on human immunodeficiency virus type 1 gp120 with griffithsin improves the antibody responses to both proteins in mice. [AIDS Res Hum Retroviruses. 28 \(2\): 206-14.](#)
4. Singh, S.M. *et al.* (2016) Characterization of Immune Responses to an Inactivated Avian Influenza Virus Vaccine Adjuvanted with Nanoparticles Containing CpG ODN. [Viral Immunol. Apr 14. \[Epub ahead of print\]](#)
5. Iwaszko-Simonik, A. *et al.* (2015) Expression of surface platelet receptors (CD62P and CD41/61) in horses with recurrent airway obstruction (RAO). [Vet Immunol Immunopathol. 164 \(1-2\): 87-92.](#)
6. Askari, N. *et al.* (2015) Tetracycline-regulated expression of OLIG2 gene in human dental pulp stem cells lead to mouse sciatic nerve regeneration upon transplantation. [Neuroscience. 305: 197-208.](#)
7. Topoluk, N. *et al.* (2017) Amniotic Mesenchymal Stromal Cells Exhibit Preferential Osteogenic and Chondrogenic Differentiation and Enhanced Matrix Production Compared With Adipose Mesenchymal Stromal Cells. [Am J Sports Med. : 363546517706138.](#)
8. Alimolaei, M. *et al.* (2017) A Recombinant Probiotic, *Lactobacillus casei*, Expressing the *Clostridium perfringens*  $\alpha$ -toxoid, as an Orally Vaccine Candidate Against Gas Gangrene and Necrotic Enteritis. [Probiotics Antimicrob Proteins. Apr 11 \[Epub ahead of print\].](#)
9. Schmidli, M.R. *et al.* (2018) Inflammatory pattern of the infrapatellar fat pad in dogs with canine cruciate ligament disease. [BMC Vet Res. 14 \(1\): 161.](#)
10. Li, T. *et al.* (2021) RNF167 activates mTORC1 and promotes tumorigenesis by targeting CASTOR1 for ubiquitination and degradation. [Nat Commun. 12 \(1\): 1055.](#)
11. Dicks, M.D.J. *et al.* (2022) Modular capsid decoration boosts adenovirus vaccine-induced humoral and cellular immunity against SARS-CoV-2 [bioRxiv: Feb 22. \[Epub ahead of print\].](#)

---

## Storage

Store at +4°C.

DO NOT FREEZE

This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10089 available at:  
<https://www.bio-rad-antibodies.com/SDS/STAR117A>  
10089

---

**Regulatory** For research purposes only

---

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M369631:200529'

**Printed on 19 Oct 2023**

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)