

Datasheet: STAR117D488GA

## **BATCH NUMBER 162837**

Description:	GOAT ANTI MOUSE IgG (H/L):DyLight®488 (MULTI SPECIES ADSORBED)
Specificity:	IgG (H/L)
Format:	DyLight®488
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

# **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/400 - 1/800
Immunofluorescence	-			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls

Target Species	Mouse		
Product Form	Purified IgG conjug	ated to DyLight <sup>®</sup> 488 - lic	duid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm
	Dylight®488	493	518
Preparation	Purified IgG prepare	ed by affinity chromatog	raphy
Antiserum Preparatio	n Antisera to mouse l purified antigen	lgG were raised by repe	ated immunisations
Buffer Solution	Phosphate buffered	d saline	
Preservative Stabilisers	0.09% Sodium Azio	de (NaN <sub>3</sub> )	
Approx. Protein	IgG concentration 1	1.0mg/ml	

### Concentrations

Immunogen	Whole mou	se IgG			
External Database	Unibert.				
Links	UniProt:	D. L. t. J			
		P01837 Related reagents			
	P01869	Related re	<del></del>		
	P01867	Related re	<del></del>		
	P01864	Related re	<del></del>		
	P01843	Related re	<del></del>		
	P01865	Related re	<del></del>		
	P01844	Related re	<del></del>		
	P01868	<del></del>			
	P01724				
	P03987 P01863	<del></del>			
	P01845				
	<u>F01043</u>	<u>rtelateu re</u>	sayents		
	Entrez Ge	ne:			
	<u>16071</u>	lgk-C	Related reagents		
	<u>16017</u>	lghg1	Related reagents		
	<u>16016</u>	lghg2b	Related reagents		
	<u>380793</u>	lgh-1a	Related reagents		
	<u>380793</u>	lgh-1a	Related reagents		
	<u>433053</u>	LOC433053	Related reagents		
	<u>16017</u>	lghg1	Related reagents		
	<u>16142</u>	lglv1	Related reagents		
	<u>110786</u>	lglc2	Related reagents		
	<u>110787</u>	Iglc3	Related reagents		
	<u>380793</u>	lgh-1a	Related reagents		
	<u>380795</u>	Al324046	Related reagents		
Synonyms	lgh-4				
RRID	AB_108518	328			
Specificity		Mouse IgG ant	<b>tibody</b> recognizes mouse IgG and light chains common to asses.		
		-	been cross-adsorbed using human, bovine, porcine, equir noabsorbants to remove cross-reactive antibodies. Less th		

Goat anti Mouse IgG has been cross-adsorbed using human, bovine, porcine, equine, lapine and chicken immunoabsorbants to remove cross-reactive antibodies. Less than 0.1% cross reactivity was detected to human, bovine, porcine, equine, caprine, lapine and chicken IgG by immunoelectrophoresis and ELISA.

Goat anti Mouse IgG antibody is highly recommended for use as a secondary antibody

with human and veterinary samples. Goat anti Mouse IgG antibody has been used successfully as a secondary detection reagent in combination with mouse clone  $\underline{\text{CC327}}$  for the detection of TNF $\alpha$  and mouse clone  $\underline{8\text{M6}}$  for the detection of interleukin-8 in bovine respiritory syncitial virus infected, neonatal ovine lung tissue by immunohistochemistry (Redondo *et al.* 2013).

#### Flow Cytometry

Use 50 ul of the suggested working dilution to label 1x10<sup>6</sup> cells in 100ul

#### References

- 1. Abdala-Valencia, H. *et al.* (2012) Vitamin E isoforms differentially regulate intercellular adhesion molecule-1 activation of PKCα in human microvascular endothelial cells. <u>PLoS</u> One. 7: e41054.
- 2. Redondo, E. *et al.* (2014) Induction of interleukin-8 and interleukin-12 in neonatal ovine lung following experimental inoculation of bovine respiratory syncytial virus. <u>J Comp</u> Pathol. 150 (4): 434-48.
- 3. Banerjee, K. *et al.* (2012) Occluding the mannose moieties on human immunodeficiency virus type 1 gp120 with griffithsin improves the antibody responses to both proteins in mice. <u>AIDS Res Hum Retroviruses. 28 (2): 206-14.</u>
- 4. Singh, S.M. *et al.* (2016) Characterization of Immune Responses to an Inactivated Avian Influenza Virus Vaccine Adjuvanted with Nanoparticles Containing CpG ODN. <u>Viral Immunol</u>. Apr 14. [Epub ahead of print]
- 5. Iwaszko-Simonik, A. *et al.* (2015) Expression of surface platelet receptors (CD62P and CD41/61) in horses with recurrent airway obstruction (RAO). <u>Vet Immunol Immunopathol.</u> 164 (1-2): 87-92.
- 6. Askari, N. *et al.* (2015) Tetracycline-regulated expression of OLIG2 gene in human dental pulp stem cells lead to mouse sciatic nerve regeneration upon transplantation. Neuroscience. 305: 197-208.
- 7. Topoluk, N. *et al.* (2017) Amniotic Mesenchymal Stromal Cells Exhibit Preferential Osteogenic and Chondrogenic Differentiation and Enhanced Matrix Production Compared With Adipose Mesenchymal Stromal Cells. <u>Am J Sports Med.</u> : 363546517706138.
- 8. Alimolaei, M. *et al.* (2017) A Recombinant Probiotic, *Lactobacillus casei*, Expressing the *Clostridium perfringens* α-toxoid, as an Orally Vaccine Candidate Against Gas Gangrene and Necrotic Enteritis. Probiotics Antimicrob Proteins. Apr 11 [Epub ahead of print].
- 9. Schmidli, M.R. *et al.* (2018) Inflammatory pattern of the infrapatellar fat pad in dogs with canine cruciate ligament disease. <u>BMC Vet Res. 14 (1): 161.</u>
- 10. Li, T. *et al.* (2021) RNF167 activates mTORC1 and promotes tumorigenesis by targeting CASTOR1 for ubiquitination and degradation. <u>Nat Commun. 12 (1): 1055.</u>
- 11. Dicks, M.D.J. *et al.* (2022) Modular capsid decoration boosts adenovirus vaccine-induced humoral and cellular immunity against SARS-CoV-2 <u>bioRxiv</u>: Feb 22. [Epub ahead of print].

#### Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch
Acknowledgements	DyLight <sup>®</sup> is a trademark of Thermo Fisher Scientific Inc. and its subsidiarie
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/STAR117D488GA">https://www.bio-rad-antibodies.com/SDS/STAR117D488GA</a> 10040
Regulatory	For research purposes only

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385180:210513'

#### Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint