

Datasheet: STAR117F

BATCH NUMBER 159615

Description:	GOAT ANTI MOUSE IgG (H/L):FITC (MULTI SPECIES ADSORBED)
Specificity:	IgG (H/L)
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/200
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
Immunofluorescence			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography		
Antiserum Preparation	Antisera to mouse IgG were raised by repeated immunisations of goats with h purified antigen.		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	0.2% Bovine Serum Albumin		

Approx. Protein Concentrations	IgG concentration 0.5 mg/ml		
Immunogen	Whole mouse IgG		
External Database Links	<div>UniProt:</div> <div><div>P01837</div><div>Related reagents</div></div> <div><div>P01869</div><div>Related reagents</div></div> <div><div>P01867</div><div>Related reagents</div></div> <div><div>P01864</div><div>Related reagents</div></div> <div><div>P01843</div><div>Related reagents</div></div> <div><div>P01865</div><div>Related reagents</div></div> <div><div>P01844</div><div>Related reagents</div></div> <div><div>P01868</div><div>Related reagents</div></div> <div><div>P01724</div><div>Related reagents</div></div> <div><div>P03987</div><div>Related reagents</div></div> <div><div>P01863</div><div>Related reagents</div></div> <div><div>P01845</div><div>Related reagents</div></div> <div>Entrez Gene:</div> <div><div>16071</div><div>Igk-C</div><div>Related reagents</div></div> <div><div>16017</div><div>Ighg1</div><div>Related reagents</div></div> <div><div>16016</div><div>Ighg2b</div><div>Related reagents</div></div> <div><div>380793</div><div>Igh-1a</div><div>Related reagents</div></div> <div><div>380793</div><div>Igh-1a</div><div>Related reagents</div></div> <div><div>433053</div><div>LOC433053</div><div>Related reagents</div></div> <div><div>16017</div><div>Ighg1</div><div>Related reagents</div></div> <div><div>16142</div><div>Iglv1</div><div>Related reagents</div></div> <div><div>110786</div><div>Iglc2</div><div>Related reagents</div></div> <div><div>110787</div><div>Iglc3</div><div>Related reagents</div></div> <div><div>380793</div><div>Igh-1a</div><div>Related reagents</div></div> <div><div>380795</div><div>AI324046</div><div>Related reagents</div></div>		
Synonyms	Igh-4		
RRID	AB_324190		
Specificity	Goat anti Mouse IgG antibody recognizes mouse IgG and light chains common to other mouse immunoglobulin classes.		

Goat anti Mouse IgG antibody is highly recommended for use as a secondary antibody with human and veterinary samples. Goat anti Mouse IgG antibody has been used successfully as a secondary detection reagent in combination with mouse clone [CC327](#) for the detection of TNF α and mouse clone [8M6](#) for the detection of interleukin-8 in bovine respiratory syncytial virus infected, neonatal ovine lung tissue by immunohistochemistry ([Redondo et al. 2013](#)).

Flow Cytometry	Use 50 ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul
-----------------------	---

References	<ol style="list-style-type: none"> 1. Abdala-Valencia, H. et al. (2012) Vitamin E isoforms differentially regulate intercellular adhesion molecule-1 activation of PKCα in human microvascular endothelial cells. PLoS One. 7: e41054. 2. Redondo, E. et al. (2014) Induction of interleukin-8 and interleukin-12 in neonatal ovine lung following experimental inoculation of bovine respiratory syncytial virus. J Comp Pathol. 150 (4): 434-48. 3. Banerjee, K. et al. (2012) Occluding the mannose moieties on human immunodeficiency virus type 1 gp120 with griffithsin improves the antibody responses to both proteins in mice. AIDS Res Hum Retroviruses. 28 (2): 206-14. 4. Singh, S.M. et al. (2016) Characterization of Immune Responses to an Inactivated Avian Influenza Virus Vaccine Adjuvanted with Nanoparticles Containing CpG ODN. Viral Immunol. Apr 14. [Epub ahead of print] 5. Iwaszko-Simonik, A. et al. (2015) Expression of surface platelet receptors (CD62P and CD41/61) in horses with recurrent airway obstruction (RAO). Vet Immunol Immunopathol. 164 (1-2): 87-92. 6. Askari, N. et al. (2015) Tetracycline-regulated expression of OLIG2 gene in human dental pulp stem cells lead to mouse sciatic nerve regeneration upon transplantation. Neuroscience. 305: 197-208. 7. Topoluk, N. et al. (2017) Amniotic Mesenchymal Stromal Cells Exhibit Preferential Osteogenic and Chondrogenic Differentiation and Enhanced Matrix Production Compared With Adipose Mesenchymal Stromal Cells. Am J Sports Med. : 363546517706138. 8. Alimolaei, M. et al. (2017) A Recombinant Probiotic, <i>Lactobacillus casei</i>, Expressing the <i>Clostridium perfringens</i> α-toxoid, as an Orally Vaccine Candidate Against Gas Gangrene and Necrotic Enteritis. Probiotics Antimicrob Proteins. Apr 11 [Epub ahead of print]. 9. Schmidli, M.R. et al. (2018) Inflammatory pattern of the infrapatellar fat pad in dogs with canine cruciate ligament disease. BMC Vet Res. 14 (1): 161. 10. Li, T. et al. (2021) RNF167 activates mTORC1 and promotes tumorigenesis by targeting CASTOR1 for ubiquitination and degradation. Nat Commun. 12 (1): 1055. 11. Dicks, M.D.J. et al. (2022) Modular capsid decoration boosts adenovirus vaccine-induced humoral and cellular immunity against SARS-CoV-2 bioRxiv: Feb 22. [Epub ahead of print].
-------------------	--

Storage	<p>Store at +4°C. DO NOT FREEZE.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected from light.</p> <p>Should this product contain a precipitate we recommend microcentrifugation before use.</p>
----------------	---

Guarantee	12 months from date of despatch
------------------	---------------------------------

**Health And Safety
Information**

Material Safety Datasheet documentation #10041 available at:
<https://www.bio-rad-antibodies.com/SDS/STAR117F10041>

Regulatory

For research purposes only

**North & South
America**

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M369638:200529'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)